arXiv:1205.1809v2 [math.SG] 4 Dec 2012

COUNTING GENUS ZERO REAL CURVES IN SYMPLECTIC MANIFOLDS

MOHAMMAD FARAJZADEH TEHRANI

ABSTRACT. There are two types of J-holomorphic spheres in a symplectic manifold invariant under an antisymplectic involution: those that have a fixed point locus and those that do not. The former are described by moduli spaces of J-holomorphic disks, which are well studied in the literature. In this paper, we first study moduli spaces describing the latter and then combine the two types of moduli spaces to get a well-defined theory of counting real curves of genus 0. We use equivariant localization to show that these invariants (unlike the disk invariants) are essentially the same for the two (standard) involutions on $\mathbb{P}^{4n-1}$. Some followup papers extends these ideas to genus 1 and higher.

CONTENTS

1. Introduction 2
1.1. Disk or τ-invariants 3
1.2. η-invariants 4
1.3. Real GW invariants 6
1.4. Projective spaces 7
1.5. acknowledgment 9
2. Moduli spaces of real curves without fixed points 10
2.1. Orientation 11
2.2. Kuranishi structure 15
3. Real GW invariants 16
Appendix A. Computations [Joint with Aleksey Zinger] 21
A.1. Odd dimensional projective space 21
A.2. The canonical orientations 25
A.3. Fixed loci 26
A.4. Normal bundles 27
A.5. Applications 29
A.6. Proof of Lemma A.2 30
References 35

Date: March 3, 2013.

1. Introduction

Let (X, ω, ϕ) be a symplectic manifold, which we will assume to be connected throughout this paper, with a real structure ϕ, i.e a diffeomorphism $\phi\colon X \to X$ such that $\phi^2 = \mathrm{id}_X$ and $\phi^*\omega = -\omega$. Let $L = \mathrm{Fix}(\phi) \subset X$ be the fixed point locus of ϕ; L is a Lagrangian submanifold of (X, ω) which can be empty. In the simplest case of $(X, \omega) = (\mathbb{P}^1, \omega_{\mathrm{FS}})$, where ω_{FS} is the Fubini-Study symplectic form, there are involutions of both types. An almost complex structure J on TX is called (ω, ϕ)-compatible if $\phi^*J = -J$ and $\omega(\cdot, J\cdot)$ is a metric. Denote the set of such almost complex structures by $\mathcal{J}_{\omega,\phi}$ or simply $\mathcal{J}_\phi$.

Fix a compatible almost complex structure J. Let $u\colon \mathbb{P}^1 \to X$ be an n-marked somewhere injective J-holomorphic sphere, i.e.

$$du + J \circ du \circ j = 0, \qquad u^{-1}(u(z)) = \{z\} \quad \text{for almost every } z \in \mathbb{P}^1,$$

where j is the complex structure of $\mathbb{P}^1$. We call such a J-holomorphic map real if its image (as a marked curve) is invariant under the action of ϕ. In this case, pulling back ϕ to $\mathbb{P}^1$, we get an involution on $\mathbb{P}^1$, which may or may not have fixed points and preserves the set of marked points. After a change of coordinates, an antisymplectic involution with fixed points can be written as

$$\tau\colon \mathbb{P}^1 \to \mathbb{P}^1, \qquad \tau([z, w]) = [\bar{w}, \bar{z}],$$

while a fixed point free involution can be written as

$$\eta\colon \mathbb{P}^1 \to \mathbb{P}^1, \qquad \eta([z, w]) = [\bar{w}, -\bar{z}].$$

For $k, l \in \mathbb{Z}^{\geq 0}$ and $A \in H_2(X)$, we define $\mathcal{M}_{k,l}(X, A)^{\phi,\tau}$ and $\mathcal{M}_l(X, A)^{\phi,\eta}$ to be the moduli spaces of degree A genus zero J-holomorphic curves $u\colon \mathbb{P}^1 \to X$ satisfying

$$u = \phi \circ u \circ \tau \qquad \text{and} \qquad u = \phi \circ u \circ \eta, \tag{1.1}$$

respectively, with l disjoint ordered conjugate pairs of marked points, along with k real (τ-fixed) marked points in the first case. By [13, Appendix C], these moduli spaces have real virtual dimension

$$\begin{aligned} \dim^{\mathrm{vir}} \mathcal{M}_{k,l}(X, A)^{\phi,\tau} &= \dim_{\mathbb{C}} X + c_1(A) + 2l + k - 3, \\ \dim^{\mathrm{vir}} \mathcal{M}_l(X, A)^{\phi,\eta} &= \dim_{\mathbb{C}} X + c_1(A) + 2l - 3. \end{aligned} \tag{1.2}$$

Every J-holomorphic map $u\colon \mathbb{P}^1 \to X$ in $\mathcal{M}_{k,l}(X, A)^{\phi,\tau}$ corresponds to two J holomorphic disks $u\colon (D^2, S^1) \to (X, L)$ with k boundary marked points and l $(\pm)$-decorated interior marked points, representing $\beta, -\phi_*\beta \in H_2(X, L)$; the j-th decoration is $(+)$ if the first point of the conjugate pair $(z_j, \tau(z_j))$ lies on the chosen disk and is $(-)$ otherwise. We define $\mathcal{M}^{\mathrm{disk}}_{k,l}(X, L, \beta)_{\mathrm{dec}}$ and $\mathcal{M}^{\mathrm{disk}}_{k,l}(X, L, \beta)$ to be the moduli space of such J-holomorphic disks with and without decorations, respectively. Let $\overline{\mathcal{M}}_l(X, A)^{\phi,\eta}$ and $\overline{\mathcal{M}}_{k,l}(X, A)^{\phi,\tau}$ be the stable map compactifications of $\mathcal{M}_l(X,A)^{\phi,\eta}$ and $\mathcal{M}_{k,l}(X,A)^{\phi,\tau}$, respectively. Let

$$\begin{aligned}
&\mathrm{ev}_i\colon \overline{\mathcal{M}}_l(X,A)^{\phi,\eta}\to X, && \mathrm{ev}_i([u,\Sigma,(z_j,\eta(z_j))_{j=1}^l])=u(z_i),\\
&\mathrm{ev}_i^B\colon \overline{\mathcal{M}}_{k,l}(X,A)^{\phi,\tau}\to L, && \mathrm{ev}_i^B([u,\Sigma,(w_j)_{j=1}^k,(z_j,\tau(z_j))_{j=1}^l])=u(w_i),\\
&\mathrm{ev}_i\colon \overline{\mathcal{M}}_{k,l}(X,A)^{\phi,\tau}\to X, && \mathrm{ev}_i([u,\Sigma,(w_j)_{j=1}^k,(z_j,\tau(z_j))_{j=1}^l])=u(z_i),
\end{aligned}\tag{1.3}$$

be the natural evaluation maps.

For the classic moduli space $\overline{\mathcal{M}}_n(X,A)$ of J-holomorphic spheres in a homology class A, **Gromov-Witten invariants** are defined via integrals of the form

$$\langle\theta_1,\cdots,\theta_n\rangle_A=\int_{[\overline{\mathcal{M}}_n(X,A)]^{\mathrm{vir}}}\mathrm{ev}_1^*(\theta_1)\wedge\cdots\wedge\mathrm{ev}_n^*(\theta_n),\tag{1.4}$$

where θ_i's are cohomology classes on X; see [6],[12],[17]. These integrals make sense and are independent of J, because $\overline{\mathcal{M}}_n(X,A)$ has a virtually orientable fundamental cycle without real codimension one boundary. One would like to define similar invariants for the moduli spaces $\overline{\mathcal{M}}_{k,l}^{\mathrm{disk}}(X,L,\beta)$ and the evaluation maps in (1.3). The existence of such invariants is predicted by physicists ([2], [11], [14], [21]), but there are obstacles to defining such invariants mathematically. In addition to the transversality issues (which are also present in the classical case), issues concerning orientability and codimension one boundary arise.

1.1. Disk or τ-invariants.

Whereas moduli spaces of closed curves have a canonical orientation induced by J, $\overline{\mathcal{M}}_{k,l}^{\mathrm{disk}}(X,L,\beta)$ is not necessarily orientable. Moreover, if it is orientable, there is no canonical orientation. If L has a spin (or relative spin) structure, then $\overline{\mathcal{M}}_{k,l}^{\mathrm{disk}}(X,L,\beta)$ is orientable and a choice of spin structure canonically determines an orientation on $\overline{\mathcal{M}}_{k,l}^{\mathrm{disk}}(X,L,\beta)$; see [7, Theorem 8.1.1].

Let $j:H_2(X)\to H_2(X,L)$ be the inclusion homomorphism. The union of moduli spaces $\overline{\mathcal{M}}_{k,l}^{\mathrm{disk}}(X,L,\beta)_{\mathrm{dec}}$ over all $\beta\in H_2(X,L)$ such that $j(A)=\beta-\phi_*\beta$ is an étale double covering of $\overline{\mathcal{M}}_{k,l}(X,A)^{\phi,\tau}$, with the deck transformation

$$\tau_{\mathcal{M}}:[u,(w_j)_{j=1}^k,(z_j,\epsilon_j)_{j=1}^l]\to[\phi\circ u\circ c,(w_j)_{j=k}^1,(c(z_j),-\epsilon_j)_{j=1}^l],$$

where $\epsilon_j=\pm$ is the decoration and $c(z)=\bar z$. A choice of spin structure on $L=\mathrm{Fix}(\phi)$ determines an orientation on $\overline{\mathcal{M}}_{k,l}^{\mathrm{disk}}(X,L,\beta)_{\mathrm{dec}}$, with the anticomplex orientation imposed on the tangent spaces at the $(-)$ marked points, as in [9, Section 3.2]. By [8, Theorem 1.3] and [9, Section 3.3], $\tau_{\mathcal{M}}$ is orientation-preserving if and only if $\frac{\mu(\beta)}{2}+k$ is even, where $\mu(\beta)\in 2\mathbb{Z}$ is the Maslov index of β. In particular, if L is spin and $4|K_X$, then $\overline{\mathcal{M}}_{0,l}(X,A)^{\phi,\tau}$ or simply $\overline{\mathcal{M}}_l(X,A)^{\phi,\tau}$ is orientable, while $\tau_{\mathcal{M}}$ is orientation-reversing on $\overline{\mathcal{M}}_{1,l}^{\mathrm{disk}}(X,L,\beta)_{\mathrm{dec}}$.

The moduli spaces $\overline{\mathcal{M}}_{k,l}(X,A)^{\phi,\tau}$ have two types of codimension one boundary; see Figure 1. The first type, called **disk bubbling**, consists of maps from two spheres with a real point in common. This boundary breaks into unions of components isomorphic to

$$\mathcal{M}_{k_1+1,l_1}(X,A_1)^{\phi,\tau}\times_{(\mathrm{ev}_1^B,\mathrm{ev}_1^B)}\mathcal{M}_{k_2+1,l_2}(X,A_2)^{\phi,\tau}/G,\tag{1.5}$$



FIGURE 1. Half of the codimension one boundary curves in $\mathcal{M}_{k,l}(X,A)^{\phi,\tau}$

where

$$l_1+l_2=l, \quad k_1+k_2=k, \quad A_1+A_2=A, \quad G=\begin{cases}\mathbb{Z}_2, & \text{if } l=0,\ A_1=A_2;\\ \{1\}, & \text{otherwise.}\end{cases}$$

The second type, called sphere bubbling, appears only if $k=0$ and $A=B-\phi_*B$ for some $B\in H_2(X)$. It consists of maps from nodal domain $\Sigma=\mathbb{P}^1\cup_q\mathbb{P}^1$, taking the node q to L. This boundary is isomorphic to ($\mathbb{Z}_2$-quotient of)

$$\bigsqcup_{\substack{B\in H_2(X)\\ B-\phi_*B=A}} (\mathcal{M}_{1+l}(X,B)_{\mathrm{dec}}\times_{\mathrm{ev}_1} L), \tag{1.6}$$

where the intersection point with L, which correspond to the first marked point in the $1+l$ marked points, has no decoration. Note that the natural extension of $\tau_{\mathcal{M}}$ preserves each component of the domain of every map in the first case and interchanges them in the second case.

The boundary problem is present in nearly all cases. It has been overcome in a number of cases by either adding other terms to compensate for the effect of the boundary ([19], [20], [5]) or by gluing boundary components to each other to get moduli spaces without boundary ([18], [9]). None of these methods can address the issue of sphere bubbling; we address it in this paper.

1.2. **η-invariants.** The moduli spaces $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ have mostly been ignored in the literature. As we show, the codimension one boundary consists of maps from a wedge of two spheres taking the node to L. The restrictions of each map to the two spheres determine elements of $\mathcal{M}_{1+l}(X,B)_{\mathrm{dec}}$ and $\mathcal{M}_{1+l}(X,-\phi_*B)_{\mathrm{dec}}$ that differ by the involution

$$\begin{aligned}\phi_{\mathcal{M}}\colon \mathcal{M}_{1+l}(X,B)_{\mathrm{dec}}\times_{\mathrm{ev}_1} L &\to \mathcal{M}_{1+l}(X,-\phi_*B)_{\mathrm{dec}}\times_{\mathrm{ev}_1} L,\\ [u,z_0,(z_1,\epsilon_1),\cdots,(z_l,\epsilon_l)] &\to [\phi\circ u\circ c, c(z_0),(c(z_1),-\epsilon_1),\cdots,(c(z_l),-\epsilon_l)],\end{aligned} \tag{1.7}$$

where $c\colon \mathbb{P}^1\to\mathbb{P}^1$, and $c(z)=\bar{z}$. Thus, the codimension one boundary breaks into unions of components isomorphic to

$$\mathcal{M}_{1+l}(X,B)_{\mathrm{dec}}\times_{\mathrm{ev}_1} L, \tag{1.8}$$

with $B\in H_2(X)$ such that $A=B-\phi_*B$. In particular, if $\mathrm{Fix}(\phi)=\emptyset$, there are no codimension one boundary components, and we obtain the following result.

Proposition 1.1. *If (X,ω,ϕ) is a symplectic manifold with a real structure ϕ and $\mathrm{Fix}(\phi)=\emptyset$, $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ has a topology with respect to which it is compact and Hausdorff. It has a Kuranishi structure without boundary of virtual real dimension*

$$d=c_1(A)+\dim_{\mathbb{C}}X-3+2l.$$

Thus, it determines an element of $H_d(\overline{\mathcal{M}}_l(X,A)^{\phi,\eta},\mathcal{O})$, where $\mathcal{O}$ is the orientation bundle.

This proposition is proved in Section 2.

Remark 1.2. There are many symplectic manifolds (X,ω) admitting antisymplectic involutions without fixed points. For example, the involution η_{2m-1} on $\mathbb{P}^{2m-1}$ defined in (1.8) has no fixed points. Furthermore, the symplectic cut of [3, Section 2] associates to each real symplectic manifold (X,ω,ϕ) with $\mathrm{Fix}(\phi)\cong S^n,\mathbb{RP}^n$ a real symplectic manifold (X_+,ω_+,ϕ_+) with $\mathrm{Fix}(\phi_+)=\emptyset$ by "cutting out" $\mathrm{Fix}(\phi)$ and replacing that with a divisor.

In order to define invariants, we also need to consider the orientation problem, which has not been studied before. A **real structure** on a vector bundle $E\to X$ is an anticomplex linear involution $\phi_E\colon E\to E$ covering ϕ. A **real square root** of a complex line bundle $E\to X$ with real structure ϕ_E is a complex line bundle $E'\to X$ with real structure $\phi_{E'}$ such that

$$(E,\phi_E)\cong(E'\otimes E',\phi_{E'}\otimes\phi_{E'}).$$

The involution ϕ on X canonically lifts to an involution ϕ_{K_X} on the complex line bundle $K_X=\Lambda_{\mathbb{C}}^{\mathrm{top}}T^*X$.

Theorem 1.3. *Let (X,ω,ϕ) be a symplectic manifold with a real structure. If (K_X,ϕ_{K_X}) admits a real square root, all moduli spaces $\mathcal{M}_l(X,A)^{\phi,\eta}$ are orientable. Moreover, a choice of real isomorphism $(K_X,\phi_{K_X})\cong(E\otimes E,\phi_E\otimes\phi_E)$ canonically determines the orientation.*

This theorem is proved in Section 2.1.

Remark 1.4. If $L\to\mathbb{P}^1$ is a holomorphic line bundle with a complex antilinear lift $\tilde{\eta}$ of $\eta\colon\mathbb{P}^1\to\mathbb{P}^1$, for all $k\in\mathbb{Z}$ there is a decomposition

$$H^0(L\otimes(T\mathbb{P}^1)^{\otimes k})=H^0_+(L\otimes(T\mathbb{P}^1)^{\otimes k})\oplus H^0_-(L\otimes(T\mathbb{P}^1)^{\otimes k})$$

into the ± 1 eigenspaces of the endomorphism

$$H^0(L \otimes (T\mathbb{P}^1)^{\otimes k}) \to H^0(L \otimes (T\mathbb{P}^1)^{\otimes k}), \quad \xi \to \tilde{\eta} \circ \xi \circ \eta;$$

the two eigenspaces are interchanged by J. Since the action of η on $\mathbb{P}^1$ has no fixed points and $H^0(L \otimes (T\mathbb{P}^1)^{\otimes k})$ is nonzero for k large enough, the zeros of every element of $H^0_+(L \otimes (T\mathbb{P}^1)^{\otimes k})$ come in pairs and thus $\deg L$ is even. Hence, if $\overline{\mathcal{M}}_l(X, A)^{\phi,\eta}$ is non-empty, then $2|K_X(A)$. Thus, if K_X has a real square root, then $4|K_X(A)$ whenever $\mathcal{M}_l(X, A)^{\phi,\eta}$ is non-empty. This condition is not sufficient though.

If (X, ω, ϕ) is a Kahler manifold with a complex conjugation ϕ and $E' \to X$ is a holomorphic line bundle, then $E' \otimes \overline{\phi^* E'}$ is a holomorphic line bundle with a real structure. Hence, if $E \to X$ is a holomorphic line bundle, $E = E' \otimes E'$, and $\overline{\phi^* E'} \cong E'$, then E admits a real structure. Suppose $4|K_X$, i.e. there is a divisor D such that $K_X = [4D]$. Since $\overline{\phi^* K_X} = K_X$, it follows that $[D - \overline{\phi_* D}]$ is torsion.

Lemma 1.5. *Let (X, ω, ϕ) be a Kahler manifold with a complex conjugation ϕ. If $K_X \cong [4D]$ and $[D - \overline{\phi_* D}] = \mathcal{O}_X$, then K_X admits a real square root.*

We prove this lemma in Section 2.1. An example with $\overline{\mathcal{M}}_l(X, A)^{\phi,\eta}$ non-orientable is described in Section 2.1. However, we are not aware of any example with X simply connected and $\overline{\mathcal{M}}_l(X, A)^{\phi,\eta}$ not orientable.

1.3. **Real GW invariants.** If $L = \mathrm{Fix}(\phi) \neq \emptyset$ and the sphere bubbling is present ($k = 0$ and $A = B - \phi_* B$ for some $B \in H_2(X)$), we cannot define either the τ-invariants nor the η-invariants separately. It is noted in [15, Section 1.5], that in order to get well-defined invariants in these case, the moduli spaces $\overline{\mathcal{M}}_l(X, A)^{\phi,\tau}$ and $\overline{\mathcal{M}}_l(X, A)^{\phi,\eta}$ need to be combined somehow. This is achieved in this paper.

As described in Sections 1.1 and 1.2, the codimension one boundary corresponding to sphere bubbling in $\overline{\mathcal{M}}_l(X, A)^{\phi,\tau}$ is the same as the codimension one boundary of $\overline{\mathcal{M}}_l(X, A)^{\phi,\eta}$. By attaching $\overline{\mathcal{M}}_l(X, A)^{\phi,\tau}$ and $\overline{\mathcal{M}}_l(X, A)^{\phi,\eta}$ along their common boundary (i.e. considering all genus 0 real curves representing class A), we obtain a moduli space $\overline{\mathcal{M}}_l(X, A)^{\phi}$ whose only codimension one boundary corresponds to disk bubbling. We then use the method of [18] and observe that the codimension one boundary components of $\overline{\mathcal{M}}_l(X, A)^{\phi}$ corresponding to disk bubbling are in fact hypersurfaces and therefore $\mathcal{M}_l(X, A)^{\phi}$ does not have any codimension one boundary.

If K_X has a real square root and L is spin, the moduli spaces $\mathcal{M}_l(X, A)^{\phi,\eta}$ and $\mathcal{M}_l(X, A)^{\phi,\tau}$ are orientable. By studying the orientation along the common boundary we show that the union is also orientable.

Theorem 1.6. *If (X, ω, ϕ) is a symplectic manifold with a real structure ϕ, $\overline{\mathcal{M}}_l(X, A)^{\phi}$ has a topology with respect to which it is compact and Hausdorff. It has a Kuranishi structure without boundary of virtual real dimension*

$$d = c_1(A) + dim_{\mathbb{C}}(X) - 3 + 2l$$

and thus determines an element of $H_d(\overline{\mathcal{M}}_l(X,A)^\phi,\mathcal{O})$*, where* $\mathcal{O}$ *is the orientation bundle. If in addition* K_X *has a real square root and* L *is spin, then* $\overline{\mathcal{M}}_l(X,A)^\phi$ *is orientable and determines an element of* $H_d(\overline{\mathcal{M}}_l(X,A)^\phi,\mathbb{Q})$*.*

We prove the first part of this theorem in Section 2.2 and the second part in Section 3. We call the resulting invariants `real` GW invariants. The moduli space $\mathcal{M}_l(X,A)^\phi$ provides a framework to define real GW invariants without any restriction on the topology of the image or the involution. If $\mathcal{M}_l(X,A)^{\phi,\eta}$ or $\mathcal{M}_l(X,A)^{\phi,\tau}$ is empty, the real invariants reduce to the disk invariants or η-invariants above. If $\mathcal{M}_l(X,A)^\phi$ is not orientable, we can still consider invariants with twisted coefficients (coefficients in the orientation bundle) as in [9].

For example, if $L\cong S^3$ and X is a real symplectic Calabi-Yau threefold, then $\mathcal{M}(X,A)^\phi$ is zero-dimensional and orientable. In this case we cannot define disk invariants or η-invariants separately. Therefore, we define real GW invariants of (X,ϕ) by

$$N_A^\phi(X)=\#[\mathcal{M}(X,A)^\phi]^{\mathrm{vir}}\in\mathbb{Q}.$$

By applying the degeneration technique of [3], we prove the following statement in Section 3.

Theorem 1.7. *Let* (X,ω,ϕ) *be a real symplectic Calabi-Yau threefold. If* $L\cong S^3$*, then*

$$N_A^\phi(X)=N_A^\eta(X_+),$$

where (X_+,ω_+,ϕ_+) *is a symplectic manifold with a real structure, canonically constructed from* X*, such that* $\mathrm{Fix}(\phi_+)=\emptyset$ *and* $N_A^\eta(X_+)$ *are the corresponding* η*-invariants.*

In [3], we show that there is a natural Hamiltonian S^1-action on a neighborhood of L in X. Applying the symplectic cut and symplectic sum procedures to this action, we build a symplectic fibration $\mathcal{X}\to\Delta$ over a disk in $\mathbb{C}$, where the smooth fibers are symplectomorphic to X and the central fiber is normal crossing, $\mathcal{X}_0=X_-\cup_D X_+$. We get an induced antisymplectic involution on $\mathcal{X}$ which fixes $X_\pm$ and restricted to X_+ has no fixed points; therefore, $N_A^\eta(X_+)$ is well-defined. This shows that in a neighborhood of certain exotic almost complex structures, there is no contribution from the τ moduli spaces.

1.4. **Projective spaces.** We now discuss in some details the case $X=\mathbb{P}^n$. The involutions $\tau,\eta\colon\mathbb{P}^1\to\mathbb{P}^1$ are special cases of the antiholomorphic involutions $\tau_{2m-1},\eta_{2m-1}\colon\mathbb{P}^{2m-1}\to\mathbb{P}^{2m-1}$, where

$$\tau_{2m-1}\big([z_1,z_2,\ldots,z_{2m-1},z_{2m}]\big)=\big([\bar z_2,\bar z_1,\ldots,\bar z_{2m},\bar z_{2m-1}]\big)\tag{1.9}$$

$$\eta_{2m-1}\big([z_1,z_2,\ldots,z_{2m-1},z_{2m}]\big)=\big([-\bar z_2,\bar z_1,\ldots,-\bar z_{2m},\bar z_{2m-1}]\big).\tag{1.10}$$

In particular,

$$\overline{\mathcal{M}}_l(\mathbb{P}^{2m-1},[d])^{\eta_{2m-1}}=\overline{\mathcal{M}}_l(\mathbb{P}^{2m-1},[d])^{\eta_{2m-1},\eta},\tag{1.11}$$

ignoring the orientations.

Suppose $\phi=\tau_{2m-1},\eta_{2m-1}$ and

$$u\colon\mathbb{P}^1\to\mathbb{P}^{2m-1},\qquad [x,y]\to\big[p_1(x,y),q_1(x,y),\ldots,p_m(x,y),q_m(x,y)\big],$$

where $p_1,q_1,\ldots,p_m,q_m$ are nonzero degree d homogeneous polynomials in two variables without a common factor. If $u\circ\eta=\phi\circ u$, every common linear factor of p_i and q_i for some i must be a common factor of $p_1,q_1,\ldots,p_m,q_m$, and so there are no such factors. The commutativity condition then implies that the zeros and poles of each rational function

$$[u,v]\to p_i(u,v)/q_i(u,v)$$

are interchanged by η, d is even if $\phi=\tau_{2m-1}$, and d is odd if $\phi=\eta_{2m-1}$. Along with the explicit description of $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ at the beginning of Section 2, this implies that

$$\begin{gathered}\overline{\mathcal{M}}_l(\mathbb{P}^{2m-1},[d])^{\tau_{2m-1},\eta}=\emptyset \quad \text{if } 2\not| d,\\ \overline{\mathcal{M}}_l(\mathbb{P}^{2m-1},[d])^{\eta_{2m-1},\eta}=\emptyset,\ \overline{\mathcal{M}}_l(\mathbb{P}^{2m-1},[d])^{\eta_{2m-1}}=\emptyset \quad \text{if } 2|d.\end{gathered} \tag{1.12}$$

Since $K_{\mathbb{P}^{4m-1}}$ has a real square root and $\mathbb{RP}^{4m-1}$ is spin, $\overline{\mathcal{M}}_l(\mathbb{P}^{2m-1},[d])^{\phi}$ is orientable for $\phi=\tau_{4m-1},\eta_{4m-1}$; see Theorem 1.6. In fact, the short exact sequence of holomorphic vector bundles

$$0\to\mathbb{P}^{n-1}\times\mathbb{C}\xrightarrow{f}n\mathcal{O}_{\mathbb{P}^{n-1}}(1)\xrightarrow{h}T\mathbb{P}^{n-1}\to0 \tag{1.13}$$

over $\mathbb{P}^{n-1}$ provides a canonical real square root for $K_{\mathbb{P}^{n-1}}$ and a spin structure for $\mathbb{RP}^{n-1}$, whenever $n=4m$; see Section A.2.

For $l,t_1,\ldots,t_l\in\mathbb{Z}^+$, we can then define

$$N_d^{\phi}(t_1,\ldots,t_l)=\int_{[\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi}]}\mathrm{ev}_1^*H^{t_1}\wedge\ldots\wedge\mathrm{ev}_l^*H^{t_l}, \tag{1.14}$$

where $H\in H^2(\mathbb{P}^{4m-1},\mathbb{Z})$ is the hyperplane class.

Theorem 1.8 (joint with Aleksey Zinger)**.** *For all $m,d,l,t_1,\ldots,t_l\in\mathbb{Z}^+$,*

$$N_d^{\eta_{4m-1}}(t_1,\ldots,t_l)=-N_d^{\tau_{4m-1}}(t_1,\ldots,t_l). \tag{1.15}$$

Furthermore, these invariants vanish if $2|d$ or $2|t_k$ for some k.

We prove this theorem in Appendix A using the equivariant localization theorem of [10]. In [4] and an early version of this paper afterward, a naive version of this theorem was presented. The theorem was extended and the necessary steps for the proof were added by Aleksey Zinger. In Section A.1 and Remarks A.5, A.6, Aleksey shows that while $K_{\mathbb{P}^{4m+1}}$ does not admit a real square root, Theorem 1.8 and its proof extend to $\mathbb{P}^{4m+1}$. Remark 1.9 below explains this in more details.

If d is odd,

$$\begin{gathered}\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1}}=\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1},\tau},\\ \overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\eta_{4m-1}}=-\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\eta_{4m-1},\eta},\end{gathered} \tag{1.16}$$

by the first statement in (1.12) and by (1.11). The sign above occurs because we reverse the orientation of $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ when gluing it to $\overline{\mathcal{M}}_l(X,A)^{\phi,\tau}$ in order to make the glued moduli space orientable; this accounts for the sign in (1.15). In this case, we show that the fixed loci in $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\eta_{4m-1},\tau}$ and $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\eta_{4m-1},\eta}$ are the same, as are their normal bundles and the corresponding restrictions of the cohomology classes being integrated; this confirms (1.15) for d odd. If d is even, $N_d^{\eta_{4m-1}}(t_1,\ldots,t_l)=0$ by the first statement in (1.12). On the other hand, in this case, the fixed loci in $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1},\tau}$ and $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1},\eta}$ are the same, as are their normal bundles and the corresponding restrictions of the cohomology classes being integrated. Since the canonical orientation on $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1},\eta}$ gets flipped when this space is glued to $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1},\tau}$, the contributions to $N_d^{\tau_{4m-1}}(t_1,\ldots,t_l)$ from the fixed loci cancel in pairs; this confirms (1.15) for d even and establishes Theorem 1.8 whenever $2|d$. Whether d is odd or even, if $2|t_k$, the contributions to $N_d^{\phi}(t_1,\ldots,t_l)$ from the fixed loci in $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi,c}$, for $c=\tau,\eta$ fixed, also cancel in pairs. This establishes the remaining vanishing statement of Theorem 1.8.

In Example A.3, we show that

$$N_1^{\eta_{4m-1}}(t_1,\ldots,t_l)=1=-N_1^{\tau_{4m-1}}(t_1,\ldots,t_l)$$

whenever all $t_1,\ldots,t_l\in\mathbb{Z}^+$ are odd and $t_1+\ldots+t_l=4m-2+l$. In particular, the signed number of real lines passing through a single point in $\mathbb{P}^{4m-1}$ is ±1, as one would expect. In Example A.4, we show that

$$N_3^{\eta_3}(3,3,3)=-1=-N_3^{\tau_3}(3,3,3),$$

i.e. the signed number of real cubics passing through 3 points in $\mathbb{P}^3$ is also ±1. A similar computation shows that

$$N_5^{\eta_3}(3,3,3,3,3)=5=-N_5^{\tau_3}(3,3,3,3,3).$$

Remark 1.9. All moduli spaces $\overline{\mathcal{M}}_l(\mathbb{P}^{2m-1},[d])^{\phi,c}$ are given orientations in Section A.1. As in the $c=\tau$ case, the orientation comes from a relative spin structure on $\mathcal{M}_{0,0}^{\mathrm{disk}}(\mathbb{P}^{2m-1},[d])$, Lemma 3.1 still applies. We show directly that so does Lemma 3.3; see Section A.1. Thus, we can define the numbers $N_d^{\phi}(t_1,\ldots,t_l)$ as in (1.14), though it is unclear in what sense these numbers are invariants. By the second statement in (1.12), the numbers $N_d^{\eta_{4m+1}}(t_1,\ldots,t_l)$ still vanish for d even. As explained in Remark A.5, the contributions to the numbers $N_d^{\tau_{4m+1}}(t_1,\ldots,t_l)$ from fixed loci in $\overline{\mathcal{M}}_l(\mathbb{P}^{4m+1},[d])^{\tau_{4m+1},\tau}$ and $\overline{\mathcal{M}}_l(\mathbb{P}^{4m+1},[d])^{\tau_{4m+1},\eta}$ still cancel in pairs for d even. For d odd, there is again a correspondence between the fixed loci in $\overline{\mathcal{M}}_l(\mathbb{P}^{4m+1},[d])^{\tau_{4m+1},\tau}$ and $\overline{\mathcal{M}}_l(\mathbb{P}^{4m+1},[d])^{\eta_{4m+1},\eta}$. The identity (1.15) thus still holds; see Remark A.5.

In Section 2, we investigate the boundary and orientation problems for moduli spaces of real curves without fixed point and define η-invariants. In Section 3, we combine the orientation problem of $\mathcal{M}(X,A)^{\phi,\tau}$ and $\mathcal{M}(X,A)^{\phi,\eta}$ and prove Theorems 1.6 and 1.7. Finally, in Appendix A, we prove Theorem 1.8.

1.5. **acknowledgment.** I would like to thank my adviser, Gang Tian, for his continuous encouragement and support, and for sharing his inspiring insights. I should thank Aleksey Zinger for his patience and help with the exposition of this paper. Through the stimulus

conversations with him the presentation of theoretical part of this paper has been enhanced. I am also thankful to Penka Georgieva for many helpful discussions.

2. MODULI SPACES OF REAL CURVES WITHOUT FIXED POINTS

In this section, we study the moduli space of real curves of genus 0 without real points. As before, let

$$\eta, \tau \colon \mathbb{P}^1 \to \mathbb{P}^1, \qquad \eta(z) = \frac{-1}{\bar{z}}, \quad \tau(z) = \frac{1}{\bar{z}}. \tag{2.1}$$

Denote by G_η the set of Mobius transformations (automorphisms of $\mathbb{P}^1$), $\rho(z) = \dfrac{az+b}{cz+d}$, commuting with η. There is an exact sequence

$$\{1\} \to U(1) \to U(2) \to G_\eta \to \{1\}.$$

Therefore, $G_\eta \cong U(2)/U(1) \cong S^3$ is a compact orientable Lie group. It acts freely and transitively on the sphere bundle $S(T\mathbb{P}^1)$ of $T\mathbb{P}^1$. The orientation on $S(T\mathbb{P}^1)$ as the boundary of the disk bundle $D(T\mathbb{P}^1)$ with its complex orientation thus induces a canonical orientation on G_η.

The involution ϕ on X induces an involution $\tilde{\phi}$ on the moduli space $\mathcal{M}_{2l}(X,A)$ of all degree A $2l$-marked somewhere injective J-holomorphic spheres:

$$\tilde{\phi}\big([u, z_1, z_2, \ldots, z_{2n-1}, z_{2l}]\big) = [\phi \circ u \circ \eta, \eta(z_2), \eta(z_1), \ldots, \eta(z_{2n}), \eta(z_{2l-1})].$$

For every J-holomorphic sphere $u \colon \mathbb{P}^1 \to X$ in the fixed point locus of $\tilde{\phi}$, there exists at most one antiholomorphic involution η_u such that $\mathrm{Fix}(\eta_u) = \emptyset$ and $u = \phi \circ u \circ \eta_u$; therefore, the fixed point locus of $\tilde{\phi}$ is precisely $\mathcal{M}_l(X,A)^{\phi,\eta}$. Intuitively, $\mathcal{M}_l(X,A)^{\phi,\eta}$ has half the dimension of $\mathcal{M}_{2l}(X,A)$.

Let $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ denote the stable map compactification of $\mathcal{M}_l(X,A)^{\phi,\eta}$. This is a closed subset of $\overline{\mathcal{M}}_{2l}(X,A)$ consisting of maps $[u, z_1, \ldots, z_{2l}]$ with the property that there exists an antiholomorphic involution η_u on the domain Σ_u of u such that

$$|\mathrm{Fix}(\eta_u)| \le 1, \qquad u = \phi \circ u \circ \eta_u, \qquad \phi_u(z_2) = z_1, \ \ldots, \ \phi_u(z_{2l}) = z_{2l-1}.$$

Thus, there are two possible cases for $\eta_u \colon \Sigma_u \to \Sigma_u$:

1. $\Sigma = \Sigma_0 \cup \bigcup_i (\Sigma_i \sqcup \Sigma_{\bar{i}})$, $\eta_u \colon \Sigma_0 \to \Sigma_0$ is an antiholomorphic involution without fixed points, and $\eta_u \colon \Sigma_i \to \Sigma_{\bar{i}}$ is an antiholomorphic map with inverse $\eta_u \colon \Sigma_{\bar{i}} \to \Sigma_i$;
2. $\Sigma = \bigcup_i (\Sigma_i \cup \Sigma_{\bar{i}})$, $\eta_u \colon \Sigma_i \to \Sigma_{\bar{i}}$ is an antiholomorphic map with inverse $\eta_u \colon \Sigma_{\bar{i}} \to \Sigma_i$.

In the second case, η_u fixes a node of Σ_u, which must be mapped by u to $\mathrm{Fix}(\phi)$, $\overline{\mathcal{M}}_l(X,A)$ contains no such elements if $\mathrm{Fix}(\phi) = \emptyset$.

The virtual codimension of a boundary stratum of $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ is the number of nodes in the domains of the elements of the stratum. If $\mathrm{Fix}(\phi) = \emptyset$, $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ contains no elements of the second type above, and so its boundary strata have codimension at least two. Thus, $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ is a moduli space without codimension one boundary if $\mathrm{Fix}(\phi) = \emptyset$, and there is a hope of defining GW-type invariants directly from $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$.

We study the orientation problem for $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ in Section 2.1 and describe a Kuranishi structure in Section 2.2.

2.1. **Orientation.**

In the orientation problem for $\mathcal{M}_l(X,A)^{\phi,\eta}$, it is sufficient to consider the case $l=0$ because any pair of marked points $(z_i,\overline{z_i})$ increases the tangent space by $T_{z_i}\mathbb{P}^1$, which has a canonical orientation. Let $\mathcal{P}_0(X,A)^{\phi,\eta}$ be the moduli space of parametrized J-holomorphic maps so that

$$\mathcal{M}_0(X,A)^{\phi,\eta}=\mathcal{P}_0(X,A)^{\phi,\eta}/G_\eta.$$

In order to put an orientation on $\mathcal{M}_0(X,A)^{\phi,\eta}$, it is enough to orient $\mathcal{P}_0(X,A)^{\phi,\eta}$. For this, we need to orient the determinant of the index bundle

$$\Lambda^{\mathrm{top}}H^0(E)_{\mathbb{R}}\otimes\Lambda^{\mathrm{top}}(H^1(E)_{\mathbb{R}})^*,$$

where $E=u^*TX$ and $H^0(E)_{\mathbb{R}}$ and $H^1(E)_{\mathbb{R}}$ are the kernel and cokernel of a real Cauchy-Riemann operator on E. Recall that E admits an anticomplex linear involution T_ϕ; see the left diagram in (2.7).

Definition 2.1. Let $E\to\mathbb{P}^1$ be a complex vector bundle with a real structure ϕ covering η. We call a trivialization of E over $\mathbb{C}^*$,

$$\begin{tikzcd}
E \arrow[r, "\psi"] \arrow[d, "\pi"'] & \mathbb{C}^*\times\mathbb{C}^m \arrow[d, "\pi"'] \\
\mathbb{C}^* \arrow[r, "\mathrm{id}"] & \mathbb{C}^*
\end{tikzcd}$$

admissible if the involution $\phi_\psi(z)=\psi_{\eta(z)}\circ\phi\circ\psi_z^{-1}$ coincides with the standard involution $C\colon(z,v)\to(\eta(z),\bar{v})$. Admissible trivializations ψ and ψ' of (E,ϕ) over $\mathbb{C}^*$ are called homotopic if there is a family of such trivializations ψ_t, $t\in[0,1]$, such that $\psi_0=\psi$ and $\psi_1=\psi'$.

Lemma 2.2. *For every complex vector bundle $E\to\mathbb{P}^1$ with a real structure ϕ covering η, there are two homotopy classes of admissible trivializations over $\mathbb{C}^*$. Moreover, for every admissible trivialization ψ and every map*

$$R_{(\epsilon_i)}\colon\mathbb{C}^*\times\mathbb{C}^m\to\mathbb{C}^*\times\mathbb{C}^m,\quad R_{(\epsilon_i)}(z,v)=(\epsilon_1v_1,\cdots,\epsilon_mv_m),\quad \epsilon_i=\pm1,$$

$R_{(\epsilon_i)}\circ\psi$ is another admissible trivialization which is in same homotopy class as ψ if and only if $\prod\epsilon_i=1$.

Proof. (1) As a complex vector bundle, E is trivial over $\mathbb{C}^*$. Therefore, we can fix a trivialization $\psi\colon E\to\mathbb{C}^*\times\mathbb{C}^m$. The involution ϕ then corresponds to a map

$$\phi_\psi\colon\mathbb{C}^*\to\mathrm{GL}(2m,\mathbb{R})$$

whose image lies in the set of anticomplex linear matrices. We find a change of trivialization matrix

$$A\colon\mathbb{C}^*\to\mathrm{GL}(m,\mathbb{C})\qquad\text{s.t.}\quad A_{\eta(z)}\circ\phi_\psi\circ A_z^{-1}=C. \tag{2.2}$$

Let $B_\psi(z) = C \circ \phi_\psi(z) \in \mathrm{GL}(m,\mathbb{C})$. Since ϕ_ψ is an involution, $\overline{B_\psi(\eta(z))}B_\psi(z) = \mathbb{I}_m$. Composing on the left by C, we can rewrite (2.2) as

$$\overline{A_{\eta(z)}} \circ B_\psi \circ A_z^{-1} = \mathbb{I}_m. \tag{2.3}$$

Let $\alpha\colon \mathbb{H}\setminus\{0\} \to \mathrm{GL}(m,\mathbb{C})$, where $\mathbb{H}$ is the closed upper half-plane, be a family of matrices such that

$$\alpha(r) = \begin{cases} \mathbb{I}_m & \text{if } r \in \mathbb{R}^+; \\ \overline{B_\psi(\eta(r))} & \text{if } r \in \mathbb{R}^-. \end{cases}$$

Next define

$$A(z) = \begin{cases} \alpha(z)B_\psi(z) & \text{if } z \in \mathbb{H}\setminus\{0\}; \\ \overline{\alpha(\eta(z))} & \text{if } z \in \overline{\mathbb{H}}\setminus\{0\}. \end{cases}$$

It is easy to check that A is continuous and satisfies (2.3).

(2) If ψ is an admissible trivialization, any other admissible trivialization is of the form $A\circ\psi$, where

$$A\colon \mathbb{C}^* \to \mathrm{GL}(m,\mathbb{C}) \qquad \text{and} \qquad \overline{A(\eta(z))}A(z)^{-1} = \mathbb{I}_m. \tag{2.4}$$

The question is whether A is homotopic to identity through a family A_t of matrices satisfying the same equation as (2.4).

Let

$$G = \left\{\gamma\colon [0,1] \to \mathrm{GL}(m,\mathbb{C}) \mid \gamma(0) = \overline{\gamma(1)}\right\}, \quad G_0 = \{\gamma \in G : \gamma(0) = \mathbb{I}_m\};$$

the set G is a group under pointwise multiplication, while G_0 is its subgroup. The restriction of A to the upper semi-circle, $\{z = e^{i\pi t} \mid t \in [0,1]\}$, determines an element of G. In fact, the space of A's satisfying (2.3) is homotopic to G. The map

$$G \to \mathrm{GL}(m,\mathbb{C}), \quad \gamma \to \gamma(0),$$

is a fiber bundle with fiber G_0. From the associated long exact sequence,

$$\cdots \to \pi_1(\mathrm{GL}(m,\mathbb{C})) \to \pi_0(G_0) \to \pi_0(G) \to \pi_0(\mathrm{GL}(m,\mathbb{C})) \to 0,$$

we conclude that $\pi_0(G) = \mathbb{Z}/2\mathbb{Z}$.[1] Therefore, there are two homotopy classes of admissible trivializations.

The remaining claim of the lemma is checked by chasing the maps in the long exact sequence. □

Lemma 2.3. *Let $E \to \mathbb{P}^1$ be a complex vector bundle with a real structure ϕ lifting η. Every admissible trivialization of (E,ϕ) over $\mathbb{C}^* \subset \mathbb{P}^1$ canonically determines an orientation of $\Lambda^{\mathrm{top}}H^0(E)_\mathbb{R} \otimes \Lambda^{\mathrm{top}}(H^1(E)_\mathbb{R})^*$. The two orientations given by two different admissible trivializations coincide if and only if they are in the same homotopy class.*

---

[1] The homomorphism $\pi_1(\mathrm{GL}(m,\mathbb{C})) \to \pi_0(G_0) \cong \pi_1(\mathrm{GL}(m,\mathbb{C})) \cong \mathbb{Z}$ is multiplication by -2 for the following reason. The loop $\gamma : [0,1] \to \mathrm{GL}(m,\mathbb{C})$ of the diagonal matrices with the first entry $e^{2i\pi t}$ and the remaining entries 1 generates $\pi_1(\mathrm{GL}(m,\mathbb{C}))$. It lifts to the path $s \to \gamma_s$ in G given by $\gamma_s(t) = \gamma((1-2s)t)$; the end point of this path is an element of $\pi_1(\mathrm{GL}(m,\mathbb{C}))$ homotopic to -2γ.

Proof. The proof is analogous to that of [7, Proposition 8.1.4]. Contracting each of the two circles

$$C_{0,r} = \{z \in \mathbb{C}^* \mid |z| = r\} \quad \text{and} \quad C_{\infty,r} = \left\{z \in \mathbb{C}^* \mid |z| = \frac{1}{r}\right\}$$

to a point, we obtain a nodal curve $\Sigma = \Sigma_{\text{top}} \cup \Sigma_0 \cup \Sigma_{\text{bot}}$ (picture below) with an induced fixed point free involution η_Σ. We denote the quotient map by $\pi\colon \mathbb{P}^1 \to \Sigma$. Denote by q and $\eta_\Sigma(q)$ the nodal points of Σ. We may assume that q and $\eta_\Sigma(q)$ are respectively 0 and ∞ in $\Sigma_0 \cong \mathbb{P}^1$.



Via the given trivialization, the bundle (E, ϕ) descends to a bundle $(\tilde{E}, \tilde{\phi})$ over Σ so that

$$\tilde{E}\mid_{\Sigma_0} \cong \mathbb{P}^1 \times \mathbb{C}^m$$

and the involution $\tilde{\phi}\mid_{\Sigma_0}$ sends (z, v) to $(\eta_\Sigma(z), \bar{v})$. Over $\Sigma_{\text{top}} \cup \Sigma_{\text{bot}}$, $\tilde{\phi}$ is an anticomplex linear map of the form

$$\tilde{\phi}\colon \tilde{E}\mid_{\Sigma_{\text{top}}} \to \tilde{E}\mid_{\Sigma_{\text{bot}}}.$$

A section of $(\tilde{E}, \tilde{\phi})$ is of the form $\xi = (\xi_{\text{top}}, \xi_0, \xi_{\text{bot}})$, with matching conditions at the nodes. A section ξ is real if and only if

$$\xi_{\text{bot}}(\eta_\Sigma(z)) = \tilde{\phi}(\xi_{\text{top}}(z)), \forall z \in \Sigma_{\text{top}} \quad \text{and} \quad \xi_0 \in \Gamma(\tilde{E}\mid_{\Sigma_0})_{\mathbb{R}}.$$

Therefore, it is determined by an arbitrary section of $\tilde{E}\mid_{\Sigma_{\text{top}}}$ and a real section of $\tilde{E}\mid_{\Sigma_0}$ which match at q.

The matching condition at the nodes gives a short exact sequence

$$0 \to W^{1,p}(\tilde{E})_{\mathbb{R}} \to W^{1,p}(\tilde{E}\mid_{\Sigma_{\text{top}}}) \oplus W^{1,p}(\tilde{E}\mid_{\Sigma_0})_{\mathbb{R}} \to \mathbb{C}^m_q \to 0.$$

The associated index of the pair $(\tilde{E}, \tilde{\phi})$ is given by

$$\text{ind}_{\mathbb{R}}\tilde{E} \cong \text{ind}_{\mathbb{C}}(\tilde{E}\mid_{\Sigma_{\text{top}}}) \otimes \text{index}_{\mathbb{R}}(\tilde{E}\mid_{\Sigma_0}) \otimes \det{}_{\mathbb{R}}(\mathbb{C}^m_q)^*. \tag{2.5}$$

Over Σ_0, the index bundle is canonically isomorphic (after deforming the Cauchy-Riemann operator) to

$$\Lambda^{\text{top}} H^0(\mathbb{P}^1 \times \mathbb{C}^m)_{\mathbb{R}} = \Lambda^{\text{top}}_{\mathbb{R}} \mathbb{R}^m \subset \Lambda^{\text{top}}_{\mathbb{C}} \mathbb{C}^m.$$

It inherits an orientation from the choice of trivialization. Since $\text{ind}_{\mathbb{C}}(\tilde{E}\mid_{\Sigma_{\text{top}}})$ and $\det_{\mathbb{R}}(\mathbb{C}^m_q)^*$ carry orientations induced by their complex structures, they are canonically oriented. Thus, (2.5) induces an orientation on $\text{ind}_{\mathbb{R}}\tilde{E}$. □

By Lemma 2.3, a systematic way of trivializing u^*TX over $\mathbb{C}^* \subset \mathbb{P}^1$ would orient $\mathcal{P}_0(X, A)^{\phi,\eta}$. Let $K_X = \Lambda^{\text{top}}_{\mathbb{C}} T^*X$ be the canonical complex line bundle over X. It inherits an involution $K_\phi\colon K_X \to K_X$ (covering ϕ) from T_ϕ. Therefore, it is a complex line bundle with an involution. Any admissible trivialization of $u^*TX\mid_{\mathbb{C}^*}$ canonically induces an admissible trivialization of $u^*K_X\mid_{\mathbb{C}^*}$ and changing the homotopy class of admissible trivialization of the former changes the homotopy class of the induced admissible trivialization. We can therefore reduce the orientation problem to the problem of finding a canonical way of admissibly trivializing u^*K_X. This is an easier problem because K_X is just a line bundle and has less structure than TX.

Let $(L,\phi_L)\to(X,\phi)$ be any complex line bundle over X with an anticomplex linear involution ϕ_L covering ϕ. The line bundle $L^{\otimes 2}$ inherits an involution from the one on L by

$$\phi_{L^{\otimes 2}}(v_1\otimes v_2)=\phi_L(v_1)\otimes\phi_L(v_2).$$

Every admissible trivialization of $u^*L\mid_{\mathbb{C}^*}$ induces an admissible trivialization of $u^*L^{\otimes 2}\mid_{\mathbb{C}^*}$. However, changing the homotopy class of trivialization of L does not change the homotopy class of the induced trivialization on $L^{\otimes 2}$, since changing the trivialization of L by the complex linear map R_{-1} of Lemma 2.2 changes the homotopy class of admissible trivialization of $L^{\otimes 2}$ by $R_{-1}\otimes R_{-1}=\mathrm{id}$. Thus, for the complex line bundle $(L^{\otimes 2},\phi_L\otimes_{\mathbb{C}}\phi_L)$ as above $u^*L^{\otimes 2}$ has a canonical admissible trivialization.

Proof of Theorem 1.3. If K_X has a real square root, then there is a natural choice of admissible trivialization, determined by the choice of $(K_X,K_\phi)\cong(L^{\otimes 2},\phi_L\otimes_{\mathbb{C}}\phi_L)$. Therefore, $\mathcal{P}_l(X,A)^{\phi,\eta}$ and hence $\mathcal{M}_l(X,A)^{\phi,\eta}$ are orientable. $\square$

proof of Lemma 1.5. Let $K_X=[4D]$, where $[\overline{\phi_*D}]=[D]$; therefore, $L:=[2D]\cong[D+\overline{\phi_*D}]$ has a real structure. Then $K_X\cong L^{\otimes 2}$ and we want to choose this isomorphism such that it commutes with ϕ_*. Let $\{U_a\}$ be a chart system symmetric with respect to ϕ, i.e. such that $\{U_{\bar a}=\phi(U_a)\}=\{U_a\}$ and $\phi:U_a\to U_{\bar a}$ is pointwise complex conjugation. Choose the trivialization of K_X given by $k_a=dz_1\wedge\cdots\wedge dz_n$ on U_a. Fix a set of defining equations $\{f_a\}$ for D on $\{U_a\}$. Then the transition maps of K_X and $[D]$ are given by $k_{ab}=k_a/k_b$ and $f_{ab}=f_a/f_b$, respectively. Define $h_a=f_a\cdot\overline{f_{\bar a}\circ\phi}$; these are local defining equations for $D+\overline{\phi_*D}$. Define h_{ab} similarly. With the above choices for the local trivializations of K_X and L, the action of the involution on local sections is given by $s_a\to s_{\bar a}=\overline{s_a\circ\phi}$. Since $K_X\cong L^{\otimes 2}$, there are $\mathbb{C}^*$-valued functions ρ_a describing the isomorphism between the two local trivializations. This isomorphism respects the involution if and only if $\rho_a=\tilde\rho_a:=\overline{\rho_{\bar a}\circ\phi}$. Define $\rho_{ab}=\rho_b/\rho_a$ and $t_a=\rho_a/\tilde\rho_a$. Since $k_{\bar a\bar b}=\overline{k_{ab}\circ\phi}$ and $h_{\bar a\bar b}=\overline{h_{ab}\circ\phi}$, it follows that $\rho_{ab}=\overline{\rho_{\bar a\bar b}\circ\phi}$. From $\rho_{ab}=\overline{\rho_{\bar a\bar b}\circ\phi}$ we conclude that $t_a=t_b$; therefore, $\{t_a\}$ represents a constant function c. Changing a with $\bar a$, we conclude $c=e^{\mathrm{i}\theta}$. Replacing ρ_a with $e^{\mathrm{i}\theta/2}\rho_a$, we get the desired isomorphism. $\square$

In Appendix A we consider $(\mathbb{P}^{4m-1},\eta_{4m-1})$; since $4|K_{\mathbb{P}^{4m-1}}$, it has a real square root. If K_X is trivial as a complex line bundle (i.e X is a symplectic Calabi-Yau manifold), then K_X has a real square root; moreover, in this case we can fix an admissible trivialization of K_X itself over X (independent of any map u) and thus determine an orientation of moduli space $\mathcal{M}_l(X,A)^{\phi,\eta}$.

As illustrated by the two examples below, there are cases where the index bundle is not orientable. The first example is similar to the non-orientable example of [7, Section 8.1.2].

Example 2.4. Let $E = S^1 \times \mathbb{P}^1 \times \mathbb{C} \to S^1 \times \mathbb{P}^1$. Define a family of involutions,

$$\phi_s \colon E\,|_{\{s\}\times\mathbb{P}^1} \to E\,|_{\{s\}\times\mathbb{P}^1}, \quad \phi_s(z,v) = (\eta(z), \overline{e^{2\pi i s}v}) \quad \forall s \in S^1.$$

The real line bundle $F \to S^1$ given by $F_s = H^0(E\,|_{\{s\}\times\mathbb{P}^1})_{\mathbb{R}}$ is then not orientable.

Example 2.5. Let $X = \mathbb{R}^2/\mathbb{Z}^2 \times \mathbb{P}^1 \times \mathbb{P}^1$, $A = \{\mathrm{pt}\} \times \mathbb{P}^1 \times \{\mathrm{pt}\} \in H_2(X)$,

$$\begin{aligned} &\phi\colon X \to X, \qquad \phi(s,t,z,w) = (s, -t, -\frac{1}{\bar{z}}, \overline{e^{2\pi i s}w}), \\ &Y = \{(s,t,w) \in \mathbb{R}^2/\mathbb{Z}^2 \times \mathbb{P}^1 : (-t, \overline{e^{2\pi i s}w}) = (t,w)\}. \end{aligned} \tag{2.6}$$

The space Y is a union of two Klein bottles with double cover

$$\mathbb{R}\cup\{\infty\} \times \mathbb{R}/\mathbb{Z} \times \{0, 1/2\} \to Y, \quad (a,s,t) \to (2s, t, ae^{2\pi i s}).$$

Let $\pi\colon X \to \mathbb{R}^2/\mathbb{Z}^2 \times \mathbb{P}^1$ be the projection to the first and third factors. Since

$$f : \mathcal{M}(X,A)^{\phi,\eta} \to Y, \qquad [u] \to \pi(\mathrm{Im}(u)),$$

is well-defined and is a diffeomorphism, it follows that $\mathcal{M}(X,A)^{\phi,\eta}$ is not orientable. If $\gamma \subset \mathcal{M}(X,A)^{\phi,\eta}$ is the preimage of the map $S^1 \to Y$, $s \to (s,0,0)$,

$$\gamma^*\mathrm{det}(T\mathcal{M}(X,A)^{\phi,\eta}) = \Lambda^{\mathrm{top}} H^0_{\mathbb{R}}(\gamma^*TX) \otimes (\Lambda^{\mathrm{top}}\mathrm{Lie}(G_\eta))^* = \mathbb{R} \otimes F,$$

where F is the unorientable line bundle in Example 2.4.

2.2. **Kuranishi structure.** If $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ is not an orbifold, in order to construct a virtual fundamental class, we need to put a Kuranishi structure on the moduli space. Such a construction for $\overline{\mathcal{M}}_{k,l}(X,A)^{\phi,\tau}$ is described in [18, Section 7]; we only describe the necessary adjustments. For simplicity, we ignore the marked points until the end of this construction.

For $(u, (z_i, \overline{z_i})_{i=1}^l) \in \mathcal{M}_l(X,A)^{\phi,\eta}$, let

$$E_u \equiv u^*TX \to \mathbb{P}^1, \quad E_u^{0,1} \equiv (T^*\mathbb{P}^1)^{0,1} \otimes_{\mathbb{C}} E_u.$$

There are commutative diagrams

$$\begin{tikzcd} E_u \arrow[r, "T_\phi"] \arrow[d, "\pi"'] & E_u \arrow[d, "\pi"'] \\ \mathbb{P}^1 \arrow[r, "\eta"] & \mathbb{P}^1 \end{tikzcd} \qquad \begin{tikzcd} E_u^{0,1} \arrow[r, "T_\phi^1"] \arrow[d, "\pi"'] & E_u^{0,1} \arrow[d, "\pi"'] \\ \mathbb{P}^1 \arrow[r, "\mathrm{id}"] & \mathbb{P}^1 \end{tikzcd} \tag{2.7}$$

where $T_\phi v = d\phi(v)$ and $T_\phi^1 \alpha = d\phi \circ \alpha \circ d\eta$. The deformation theory of $\mathcal{M}_0(X,A)^{\phi,\eta}$ is described by the linearization of the Cauchy-Riemann operator,

$$L_{J,u}\colon W^{k,p}(E_u) \to W^{k-1,p}(E_u^{0,1}), \quad p > 2, k \geq 1; \tag{2.8}$$

see [13, Chapter 3] for a similar situation. If ∇ is the Levi-Civita connection of the metric $\omega(\cdot, J\cdot)$, $L_{J,u}$ can be written as

$$L_{J,u}(\xi) = \frac{1}{2}(\nabla\xi + J\nabla\xi \circ j) - \frac{1}{2}J(\nabla_\xi J)\partial_J(u).$$

There is a commutative diagram

$$\begin{tikzcd} W^{k,p}(E_u) \arrow[r, "L_{J,u}"] \arrow[d, "\tilde{T}_\phi"'] & W^{k-1,p}(E_u^{0,1}) \arrow[d, "\tilde{T}^1_\phi"'] \\ W^{k,p}(E_u) \arrow[r, "L_{J,u}"] & W^{k-1,p}(E_u^{0,1}) \end{tikzcd}$$

where $\{\tilde{T}_\phi \xi\}(z) = T_\phi(\xi(\eta(z)))$ and $\{\tilde{T}^1_\phi \alpha\}(z) = T^1_\phi(\alpha(z))$. Let

$$\begin{aligned} W^{k,p}(E_u)_{\mathbb{R}} &= \{\xi \in W^{k,p}(E_u) \mid \tilde{T}_\phi(\xi) = \xi\}, \\ W^{k-1,p}(E_u^{0,1})_{\mathbb{R}} &= \{\alpha \in W^{k-1,p}(E_u^{0,1}) \mid \tilde{T}^1_\phi(\alpha) = \alpha\} \end{aligned} \tag{2.9}$$

denote the spaces of real sections. Let $H^0(E_u)_{\mathbb{R}}$ and $H^1(E_u)_{\mathbb{R}}$ be the kernel and cokernel, respectively, of the restricted operator

$$L_{J,u}\colon W^{k,p}(E_u)_{\mathbb{R}} \to W^{k-1,p}(E_u^{0,1})_{\mathbb{R}}.$$

If $H^1(E_u)_{\mathbb{R}} = 0$, then $\mathcal{M}(X,A)^{\phi,\eta}$ is a manifold near u of real dimension

$$\dim_{\mathbb{R}} H^0(E_u)_{\mathbb{R}} - \dim G_\eta = \operatorname{ind}_{\mathbb{R}}(L_{J,u}) - 3 = c_1(A) + \dim_{\mathbb{C}} X - 3; \tag{2.10}$$

see [13, Theorem C.1.10]. Each pair of conjugate marked points increases the dimension by two and we get the dimension formula (1.2).

If $H^1(E_u)_{\mathbb{R}} \neq 0$, we construct a Kuranishi chart around u. For this aim, we choose finite-dimensional complex subspaces $\mathcal{E}_u \subset W^{k,p-1}(E_u^{0,1})$ such that

(1) every $\xi \in \mathcal{E}_u$ is smooth and supported away from the boundary and marked points;
(2) $\tilde{T}^1_\phi(\mathcal{E}_u) = \mathcal{E}_u$;
(3) $L_{J,u}$ modulo $\mathcal{E}_u$ is surjective.

We then choose our Kuranishi neighborhood to be $V(u) = [\bar{\partial}^{-1}(\mathcal{E}_u)]_{\mathbb{R}}$ (modulo G_η), which is a smooth manifold of dimension

$$c_1(A) + \dim_{\mathbb{C}} X - 3 + 2l + \dim_{\mathbb{C}}(\mathcal{E}_u).$$

The obstruction bundle $\mathcal{E}(u)$ at each $f \in V(u)$ is obtained by parallel translation of $\mathcal{E}_u$ with respect to the induced metric of J. We thus get a Kuranishi neighborhood $(V(u), \mathcal{E}(u))$. The Kuranishi map in this case is just the Cauchy-Riemann operator $f \to \bar{\partial}(f)$.

In order to construct Kuranishi charts for u in the boundary strata of $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$, we need gluing theorems as in [7, Chapter 7]. The gluing theorems are identical to those for J-holomorphic disks; we thus omit the details and refer the reader to [7].

3. REAL GW INVARIANTS

If $L = \mathrm{Fix}(\phi)$ is non-empty, the codimension one boundary of $\overline{\mathcal{M}}_l(X,A)^{\phi,\eta}$ might be non-empty; see (1.8). An element of codimension one boundary is of the form $(u, \Sigma = \Sigma_1 \cup_q \Sigma_2)$, where $\Sigma_i = \mathbb{P}^1$, $\eta\colon \Sigma_1 \to \Sigma_2$, and $u(q) \in L$. After a suitable reparametrization, we may assume $q = 0 \in \mathbb{P}^1$ and $\eta(z) = \overline{w}$. For real parameters $\epsilon \neq 0$, we can glue Σ into a family of smooth curves

$$\Sigma_\epsilon = \{(z, w) \in \mathbb{C}^2 : zw = \epsilon\}.$$

For $\epsilon \in \mathbb{R}$, Σ_ϵ inherits a complex conjugation from η:

$$\eta_\epsilon \colon \Sigma_\epsilon \to \Sigma_\epsilon, \qquad \eta_\epsilon(z, w) = (\overline{w}, \overline{z}).$$

The fixed point set of η_ϵ is S^1 if $\epsilon > 0$ and is empty if $\epsilon < 0$. By smoothing in one direction (ϵ negative), we get real curves without fixed points in $\overline{\mathcal{M}}_l(X, A)^{\phi,\eta}$; by smoothing in the other direction (ϵ positive), we get real curves with fixed points in $\overline{\mathcal{M}}_l(X, A)^{\phi,\tau}$. We identify the common boundary and glue the two moduli spaces to get a new moduli space whose only boundary component comes from the disk bubbling. We define $\overline{\mathcal{M}}(X, A)^\phi$ to be the resulted space.

Each nodal curve in (1.5) is of the form $(u, \Sigma = \Sigma_1 \cup_q \Sigma_2)$, where $\Sigma_i = \mathbb{P}^1$, $\tau \colon \Sigma_i \to \Sigma_i$, and $q \in \mathrm{Fix}(\tau|_{\Sigma_i}) \cong S^1$. After a suitable reparametrization, we may assume $q = (z_i = 0) \in \mathbb{C} \subset \mathbb{P}^1$ and $\tau|_{\mathbb{C}} = c$, where $c(z_i) = \bar{z}_i$. For real $\epsilon \neq 0$, we can glue Σ into a family of smooth curves

$$\Sigma_\epsilon = \{(z_1, z_2) \in \mathbb{C}^2 \ : \ z_1 z_2 = \epsilon\}.$$

For $\epsilon \in \mathbb{R}$, Σ_ϵ inherits a complex conjugation from τ:

$$\tau_\epsilon \colon \Sigma_\epsilon \to \Sigma_\epsilon, \qquad \tau_\epsilon(z_1, z_2) = (\bar{z}_1, \bar{z}_2).$$

The fixed point set of τ_ϵ is S^1. By the stability condition, for each component i, either $l_i \neq 0$ or the map $u_i \equiv u|_{\Sigma_i}$ is non-trivial. If $l_i \neq 0$, we fix one of the marked points; if u_i is non-trivial and somewhere injective, we fix a somewhere injective point of the corresponding domain. By tracking the image of the chosen point, we see that gluing the map in positive and negative directions produce different J-holomorphic curves. If u_i is multiple cover and $l_i = 0$, then the obstruction bundle near $u_i \in \mathcal{M}_{1,0}(X, A_i)^{\phi,\tau}$ is non-trivial and a Kuranishi neighborhood depends on the choice of $\mathcal{E}_{u_i}$ of the previous section. By choosing $\mathcal{E}_{u_i}$ non-symmetric with respect to the deck transformation of the covering map, we can assure that gluing in different directions produce different maps. Therefore, the real codimension one boundary term (1.5), corresponding to $\epsilon = 0$, is indeed a hypersurface. This establishes the first part of Theorem 1.6, i.e. that $\overline{\mathcal{M}}(X, A)^\phi$ has the structure of a closed Kuranishi space; the real codimension one strata (1.5) and (1.8) are real codimension one hypersurfaces in $\overline{\mathcal{M}}(X, A)^\phi$.

The next two lemmas show that $\overline{\mathcal{M}}_l(X, A)^\phi$ is orientable if K_X has a real square root and L is spin. These assumptions imply that the spaces $\mathcal{M}_l(X, A)^{\phi,\tau}$ and $\mathcal{M}_l(X, A)^{\phi,\eta}$ are orientable; see the beginning of Section 1.1 and Theorem 1.3. Since K_X has a real square root, $4|c_1(A)$ for every $A \in H_2(X)$. Thus, the first lemma below implies that the orientation of $\mathcal{M}_l(X, A)^{\phi,\tau}$ extends across the hypersurface (1.8). The second lemma implies that the orientations of $\mathcal{M}_l(X, A)^{\phi,\tau}$ and $\mathcal{M}_l(X, A)^{\phi,\eta}$ can be extended across the common boundary (1.5), possibly after flipping the orientation of the latter. In Lemma 3.3, we consider the induced orientation on the boundary ∂M of an oriented manifold M to be the one given by the outward normal vector field; i.e. $TM|_{\partial M} = T\partial M \times v_{\mathrm{out}}$, as oriented vector spaces.

Lemma 3.1. *Let (X, ω, ϕ) be a symplectic manifold with a real structure such that $L = \mathrm{Fix}(\phi)$ is spin. If $4|c_1(A_i)$, then the gluing maps*

$$\mathcal{M}_{1,l_1}(X, A_1)^{\phi,\tau} \times_{(\mathrm{ev}_1^B, \mathrm{ev}_1^B)} \mathcal{M}_{1,l_2}(X, A_2)^{\phi,\tau}/G \times \begin{cases} \mathbb{R}^{\geq 0} \\ \mathbb{R}^{\leq 0} \end{cases} \to \overline{\mathcal{M}}(X, A)^{\phi,\tau}$$

given by smoothing domain with respect to the corresponding gluing parameter ϵ, are orientation-preserving.

Remark 3.2. If (X, L) is relatively spin, as in [7], all the moduli spaces $\mathcal{M}_{k,l}^{\mathrm{disk}}(X, L, \beta)$ are still orientable. Let $[V, \sigma]$ be a relative spin structure as in [8]. Then, $\phi_*[V, \sigma]$ is another relative spin structure and is equal to $\alpha \cdot [V, \sigma]$, for some $\alpha \in H^2(X, L)$; see [8]. If

$$\alpha(\beta) \equiv \frac{\mu(\beta)}{2} \mod 2, \quad \forall \beta \in H_2(X),$$

then by Theorem 1.3 and Proposition 3.10 of [8], $\mathcal{M}_l(X, A)^{\phi,\tau}$ is orientable and the conclusion of Lemma 3.1 is still true. For example, $(\mathbb{P}^{4m+1}, \mathbb{RP}^{4m+1})$ is not spin but is relatively spin and each choice of two homotopy classes of relative spin structures satisfy the identity above.

Lemma 3.3. *Let (X, ω, ϕ) be a symplectic manifold with a real structure such that $L = \mathrm{Fix}(\phi)$ is spin and K_X has a real square root. If $A, B \in H_2(X)$ are such that $A = B - \phi_* B$, the gluing maps*

$$(\mathcal{M}_1(X, B) \times_{\mathrm{ev}_1} L) \times \mathbb{R}^+ \to \mathcal{M}(X, A)^{\phi,\tau}, \mathcal{M}(X, A)^{\phi,\eta}$$

are orientation-preserving, provided the Lagrangian on the left-hand side is oriented by the chosen spin structure of TL in the first case and by the choice of the isomorphism

$$\Lambda_{\mathbb{R}}^{\mathrm{top}}(TL) \cong \big(\Lambda_{\mathbb{C}}^{\mathrm{top}}(TL)\big)_{\mathbb{R}} = (K_X^*)_{\mathbb{R}} \approx E_{\mathbb{R}}^* \otimes E_{\mathbb{R}}^*,$$

with $(K_X, \phi_{K_X}) \cong (E \otimes E, \phi_E \otimes \phi_E)$, in the second case.

Proof. A curve in the common boundary of these two moduli spaces is of the form

$$f = [u, \Sigma = \mathbb{P}^1_{\mathrm{top}} \cup_q \mathbb{P}^1_{\mathrm{bot}}],$$

with the involution c over Σ having one fixed point, the node q. We replace each such f with the unstable map

$$\tilde{f} = [\tilde{u}, \tilde{\Sigma} = \mathbb{P}^1_{\mathrm{top}} \cup \mathbb{P}^1_0 \cup \mathbb{P}^1_{\mathrm{bot}}],$$

with $\tilde{u}$ restricting to the constant $u(q)$ over the central part $\mathbb{P}^1_0$. We can view $\tilde{f}$ as an element of $\partial\overline{\mathcal{M}}(X, A)^{\phi,\tau}$ by extending the involution to $\mathbb{P}^1_0$ via $c|_{\mathbb{P}^1_0} = \tau$ and as an element of $\partial\overline{\mathcal{M}}(X, A)^{\phi,\eta}$ by extending the involution to $\mathbb{P}^1_0$ via $c|_{\mathbb{P}^1_0} = \eta$. The real automorphism group of $\tilde{f}$ restricted to the middle component is S^1. Following the orientation and gluing argument in [7, Section 8.3] and [7, Section 7.4.1], it is easy to see that the index bundles are canonically isomorphic through the gluing procedure and the isomorphism is orientation-preserving.

In fact, for each glued map f_ϵ over the glued domain $\Sigma_\epsilon \cong \mathbb{P}^1$, in order to orient $T_{f_\epsilon}\mathcal{M}(X,A)^{\phi,c}$, we fix an admissible trivialization of f_ϵ^*TX over $\mathbb{P}^1-\{0,\infty\}$ and degenerate u_ϵ^*TX into a bundle over $\tilde{\Sigma}$, such that over the central part the induced bundle is admissibly trivial. This is exactly the case with $\tilde{u}^*TX$. Over the central part, $\tilde{u}|_{\mathbb{P}^1_0}$ is trivial and the set of real sections are orientably isomorphic to $T_{u(q)}L$. This oriented isomorphism determines the orientation of index bundle of the glued map.

Therefore, in order to prove this Lemma, it remains to compare the automorphism groups of domains before and after gluing. Let

$$G_0=\mathrm{Aut}_{\mathbb{R}}([\mathbb{P}^1_{\mathrm{top}}\cup_q\mathbb{P}^1_{\mathrm{bot}}])\cong \mathrm{Aut}(\mathbb{P}^1_{\mathrm{top}},q)\subset \mathrm{PSL}(2,\mathbb{C})$$

be the identity component of he real automorphism group of Σ. This is a real 2-dimensional complex Lie group which has a canonical orientation and

$$T_f(\mathcal{P}_1(X,B)\times_{\mathrm{ev}_1}L)\cong T_f(\mathcal{M}_1(X,B)\times_{\mathrm{ev}_1}L)\oplus T_{\mathrm{id}}G_0 \tag{3.1}$$

as oriented vector spaces, with the isomorphism obtained by splitting the short exact sequence

$$0\to T_f(\mathcal{M}_1(X,B)\times_{\mathrm{ev}_1}L)\to T_f(\mathcal{P}_1(X,B)\times_{\mathrm{ev}_1}L)\cong T_{\mathrm{id}}G_0\to 0.$$

After replacing Σ with $\tilde{\Sigma}$, the real automorphism group of the domain increases by a factor of S^1 and the gluing parameter of the domain takes values in $\mathbb{C}$. To kill the extra S^1-action in both the automorphism group and the gluing parameter, as in the statement of the lemma, we consider the gluing parameter to be positive real (absolute value of the complex one) and restrict to a real 4-dimensional section of $G_0\times S^1$, given by

$$G_0'=\{(g,e^{\mathrm{i}\theta})\subset G_0\times S^1:\ dg|_{T_0\mathbb{P}^1_{\mathrm{top}}}\in R^+e^{-\mathrm{i}\theta}\},$$

which is canonically isomorphic to G_0. Let

$$\mathcal{C}=\{(z_t,z_0,z_b,\epsilon)\in\mathbb{P}^1\times\mathbb{P}^1\times\mathbb{P}^1\times\mathbb{R}^{\geq 0}|z_tz_0=\epsilon,\, z_b\overline{c(z_0)}=\epsilon,\ \epsilon\in\mathbb{R}^{\geq 0}\}.$$

This is a real one-parameter family of genus zero real curves over $\mathbb{R}^{\geq 0}$,

$$(z_t,z_0,z_b,\epsilon)\to\epsilon,$$

with the fiber-preserving involution

$$(z_t,z_0,z_b,\epsilon)\to(\bar{z}_b,c(z_0),\bar{z}_t,\epsilon),$$

that describes a gluing of the singular real curve $\tilde{\Sigma}$ into smooth real curves.

Over $\mathcal{C}$, consider the group $G_0^{\mathcal{C}}$ generated by the following set of maps

$$\begin{aligned}
&R_r\colon(z_t,z_0,z_b,\epsilon)\to(\frac{1}{r}z_t,z_0,\frac{1}{r}z_b,\frac{1}{r}\epsilon),\quad r\in\mathbb{R}^+\text{ close to }1,\\
&R_\theta\colon(z_t,z_0,z_b,\epsilon)\to(e^{-\mathrm{i}\theta}z_t,e^{\mathrm{i}\theta}z_0,e^{\mathrm{i}\theta}z_b,\epsilon),\quad e^{\mathrm{i}\theta}\in S^1,\\
&T_a\colon(z_t,z_0,z_b,\epsilon)\to(\frac{z_t+(-1)^{|c|}\epsilon^2\bar{a}}{1+az_t},\frac{z_0+a\epsilon}{1+(-1)^{|c|}\epsilon\bar{a}z_0},\frac{z_b+(-1)^{|c|}a\epsilon^2}{1+\bar{a}z_b}),\quad a\in\mathbb{C},
\end{aligned}\tag{3.2}$$

with $|c|$ defined as in (A.2). This group extends the action of G'_0 to the whole family. Restricted to each fiber $\mathcal{C}_\epsilon$, $\epsilon \neq 0$, R_θ and T_a generate the 3-dimensional real automorphism group of the fiber, G_η or (the identity component of) G_τ, depending on c. Let

$$v_1 = \frac{d}{dr} R_r|_{r=1}, \quad v_2 = \frac{d}{d\theta} R_\theta|_{\theta=0}, \quad v_3 + \mathfrak{i} v_4 = \frac{d}{da} T_a|_{a=0}.$$

Restricted to $\mathcal{C}_0$, these vector fields form an oriented basis of $T_{\mathrm{id}} G'_0$. The restriction of v_2, v_3, v_4 to $\mathcal{C}_\epsilon$, $\epsilon \neq 0$, they form an oriented basis of $T_{\mathrm{id}} G_c$. Finally, v_1 (after some positive rescaling) is a lift of the outward normal vector field $-\frac{\partial}{\partial \epsilon}$ to the family. Since the index bundle of f and the glued maps f_ϵ are orientably isomorphic, via decomposition (3.1), we get the following isomorphism of oriented vector spaces

$$T_f(\mathcal{M}_1(X,B) \times_{\mathrm{ev}_1} L) \oplus \bigoplus_{i=1}^{4} \mathbb{R}.v_i = T_{f_\epsilon}(\mathcal{M}(X,A)^{\phi,c}) \oplus \bigoplus_{i=2}^{4} \mathbb{R}.v_i.$$

Erasing the last three terms from the equation and replacing v_1 by v_{out}, we observe that

$$T_f(\mathcal{M}_1(X,B) \times_{\mathrm{ev}_1} L) \oplus v_{\mathrm{out}} = T_{f_\epsilon}(\mathcal{M}(X,A)^{\phi,c}),$$

is an isomorphism of oriented vector spaces. This establishes the claim. □

Thus, if K_X has a real square root and L is spin, choosing the spin structure appropriately, $\overline{\mathcal{M}}_l(X,A)^\phi$ is orientable and closed. In this case, for $\theta_1, \ldots, \theta_l \in H^*(X)$, we define real GW invariants by

$$N_A^\phi(\theta_1, \ldots, \theta_l) = \int_{[\overline{\mathcal{M}}_l(X,A)^\phi]^{\mathrm{vir}}} \mathrm{ev}_1^*(\theta_1) \wedge \cdots \wedge \mathrm{ev}_l^*(\theta_l).$$

We finish this section with the proof of Theorem 1.7. By [3, Proposition 2.1], there is a symplectic fibration $\pi\colon \mathcal{X} \to \Delta$ with an induced real structure $\phi_{\mathcal{X}}$ over a disk $\Delta \subset \mathbb{C}$ such that the central fiber $\mathcal{X}_0$ is a nodal-crossing symplectic manifold with real structure $\phi_{\mathcal{X}}|_{\mathcal{X}_0}$,

$$\mathcal{X}_0 = X_- \cup_D X_+, \quad D \cong \mathbb{P}^1 \times \mathbb{P}^1, \quad \phi_\pm := \phi_{\mathcal{X}}|_{X_\pm},$$

where (X_-, ϕ_-) is symplectomorphic to real quadratic hypersurface in $\mathbb{P}^4$ given by

$$x_0^2 - \sum_{i=1}^{4} x_i^2 = 0, \qquad D = (x_0 = 0),$$

$\mathrm{Fix}(\phi_+) = \emptyset$ and $c_1(TX_+) = -D$. Moreover, fibers over Δ^* are smooth and symplectically isotopic to (X, ϕ).

Let $\mathcal{Y} = \pi^{-1}([0,1]) \subset \mathcal{X}$. Each fiber of $\mathcal{Y} \to [0,1]$ is invariant under the induced involution $\phi_{\mathcal{X}}$. Fix some compatible J on $\mathcal{X}$. For $A \in H_2(X, \mathbb{Z})$, define

$$\overline{\mathcal{M}}(\mathcal{Y}, \{J_t\}_{t\in(0,1]}, A)^{\phi_{\mathcal{X}}} = \bigcup_{t\in(0,1]} \overline{\mathcal{M}}(\mathcal{X}_t, J_t, A)^{\phi_{\mathcal{X}}}.$$

Let $\overline{\mathcal{M}}(\mathcal{Y}, \{J_t\}_{t\in I}, A)^{\phi_{\mathcal{X}}}$, where $I = [0,1]$, be the relative stable map compactification of $\overline{\mathcal{M}}(\mathcal{Y}, \{J_t\}_{t\in(0,1]}, A)^{\phi_{\mathcal{X}}}$, as in [3, Section 3.2], including maps to the fiber over zero. Every element (u,Σ) of $\overline{\mathcal{M}}(\mathcal{Y},\{J_t\}_{t\in I},A)^{\phi_{\mathcal{X}}}$ in $\mathcal{X}_0$ belongs to a fiber product of relative moduli spaces over X_- and X_+ with matching conditions on D,

$$\overline{\mathcal{M}}(X_-,D,\rho,\Gamma_-)^{\phi_-}\times_{(\mathrm{ev}_{\xi^-},\mathrm{ev}_{\xi^+})}\overline{\mathcal{M}}(X_+,D,\rho,\Gamma_+)^{\phi_+}, \tag{3.3}$$

where $\overline{\mathcal{M}}(X_-,D,\rho,\Gamma_-)^{\phi_-}$ and $\overline{\mathcal{M}}(X_+,D,\rho,\Gamma_+)^{\phi_+}$ are the relative moduli spaces of real curves, possibly with disconnected domains, with the same intersection pattern ρ; $\xi^\pm$ are contact points with D, and $\Gamma_\pm$ encodes the data corresponding to the topological types of the domain and image; see [3, Section 4] for more details on the definition.

The moduli space $\overline{\mathcal{M}}(\mathcal{Y},\{J_t\}_{t\in(0,1]},A)^{\phi_{\mathcal{X}}}$ gives a cobordism between the moduli space of real curves $\overline{\mathcal{M}}(\mathcal{X},\{J_t\}_I,A)^{\phi_{\mathcal{X}}}|_{t=1}\cong\overline{\mathcal{M}}(X,J_1,A)^{\phi}$ and the moduli space of real curves in the singular fiber. By [3, Proposition 2.1], $c_1(TX_+)=-PD(D)$. Therefore, for every curve class $B\in H_2(X_+)$ with $B\cdot D>0$,

$$\dim^{\mathrm{vir}}(\overline{\mathcal{M}}(X_+,D,\rho,\Gamma_+)^{\phi_+})\le c_1(TX_+)(B)<0,$$

whenever the image curves of type Γ_+ are in homology class B. Thus, the only term with non-trivial contribution is

$$N_A^{\phi}=\#(\overline{\mathcal{M}}(X_+,A)^{\phi_+})=\#(\overline{\mathcal{M}}(X_+,A)^{\phi_+,\eta})=N_A^{\eta}(X_+),$$

where by abuse of notation A is the image of homology class $A\subset H_2(X)$ in $H_2(X_+)$ via the natural inclusion map $H_2(X)\hookrightarrow H_2(X_+)$.

Appendix A. Computations [Joint with Aleksey Zinger]

In the rest of this paper, we use equivariant localization to prove Theorem 1.8 by summing over the fixed loci of a torus action on $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi}$. As in [15, Section 3], these loci are described by graphs with one half-edge. The contribution of the complement of the half-edge to the normal bundle of the corresponding locus is standard. Lemma A.2 below determines the key contribution of the half-edge to the normal bundle and is thus analogous to [15, Lemma 6], though our argument is completely different from [15].

We explicitly describe the canonical square root structure on $K_{\mathbb{P}^{4m-1}}$ and spin structure on $\mathbb{RP}^{4m-1}$ induced by the exact sequence (1.13) in Section A.2; these are used to define the numbers (1.14). We describe the fixed loci of a natural action of

$$\mathbb{T}\equiv(S^1)^m\equiv\big\{(\zeta_1,\ldots,\zeta_m)\in\mathbb{C}^m\colon\ |\zeta_k|=1\big\}$$

on $\overline{\mathcal{M}}_l(\mathbb{P}^{2m-1},[d])^{\phi}$ in Section A.3 and their normal bundles in Section A.4. In Section A.5, we prove Theorem 1.8 and compute some low-degree real invariants. We conclude by proving Lemma A.2 in Section A.6.

A.1. Odd dimensional projective space.

Let $m,d\in\mathbb{Z}^+$, $\phi=\tau_{2m-1},\eta_{2m-1}$, and $|\phi|$ be as in Equation A.2. The space $\mathcal{P}_0(\mathbb{P}^{2m-1},[d])^{\phi,\eta}$ consists of maps of the form

$$u\colon\mathbb{P}^1\to\mathbb{P}^{2m-1},\qquad[x,y]\to\big[p_1(x,y),q_1(x,y),\ldots,p_m(x,y),q_m(x,y)\big],$$

where $p_1, q_1, \ldots, p_m, q_m$ are degree d homogeneous polynomials in two variables without a common factor. The commutativity condition on u is equivalent to

$$p_i(x,y) = A_i \prod_{r=1}^{d} (a_{i;r}x - b_{i;r}y), \qquad q_i(x,u) = \bar{B}_i \prod_{r=1}^{d} (\bar{b}_{i;r}x + \bar{a}_{i;r}y),$$

for some $A_i, B_i \in \mathbb{C}$ and $[a_{i;r}, b_{i;r}] \in \mathbb{P}^1$ such that

$$|A_i| = |B_i| \ \forall\, i = 1, \ldots, m, \qquad [A_1, \ldots, A_m] = [B_1, \ldots, B_m] \in \mathbb{P}^{m-1},$$

i.e. $(B_1, \ldots, B_m) = \zeta(A_1, \ldots, A_m)$ for some $\zeta \in S^1 \subset \mathbb{C}$. Note that if $2 \nmid |\phi| - d$, then $\mathcal{P}_0(\mathbb{P}^{2m-1}, [d])^{\phi,\eta}$ is empty. Thus, the map

$$\begin{gathered}
(\mathrm{Sym}^d\mathbb{C})^m \times \mathbb{RP}^{2m-1} \to \mathcal{P}_0(\mathbb{P}^{2m-1}, [d])^{\phi,\eta}, \\
\big([b_{1;1}, \ldots, b_{1;d}], \ldots, [b_{m;1}, \ldots, b_{m;d}], [A_1, \ldots, A_m]\big) \to \\
\Big[A_1 \prod_{r=1}^{d} (x - b_{1;r}y), \bar{A}_1 \prod_{r=1}^{d} (\bar{b}_{1;r}x + y), \ldots, A_m \prod_{r=1}^{d} (x - b_{m;r}y), \bar{A}_m \prod_{r=1}^{d} (\bar{b}_{m;r}x + y)\Big],
\end{gathered}$$

where $\mathbb{RP}^{2m-1} \equiv (\mathbb{C}^m - \{0\})/\mathbb{R}^*$, is an isomorphism over the open subset of $\mathcal{P}_0(\mathbb{P}^{2m-1}, [d])^{\phi,\eta}$ consisting of maps u such that $u([1,0])$ does not lie in any of the coordinate subspaces of $\mathbb{P}^{2m-1}$. Since the complement of this subspace is of codimension 2, this isomorphism induces an orientation on $\mathcal{P}_0(\mathbb{P}^{2m-1}, [d])^{\phi,\eta}$ and thus on $\mathcal{M}_l(\mathbb{P}^{2m-1}, [d])^{\phi,\eta}$. We show in Remark A.5 that this is the orientation on $\mathcal{M}_l(\mathbb{P}^{2m-1}, [d])^{\phi,\eta}$ induced by the canonical real square root of $K_{\mathbb{P}^{2m-1}}$, whenever m is even.

Similar reasoning applies to the space $\mathcal{P}_0(\mathbb{P}^{2m-1}, [d])^{\tau_{2m-1},\tau}$. In this case,

$$p_i(x,y) = A_i \prod_{r=1}^{d} (a_{i;r}x + b_{i;r}y), \qquad q_i(x,u) = \bar{B}_i \prod_{r=1}^{d} (\bar{b}_{i;r}x + \bar{a}_{i;r}y)$$

with the m sets $\{[a_{i;r}, b_{i;r}] : r = 1, \ldots, d\}$ having no element of $\mathbb{RP}^1 = \mathrm{Fix}(\tau)$ in common. In particular, the map

$$\big\{(a,b) \in \mathbb{C}^2 \colon\ |a| \neq |b|\big\}/\mathbb{R}^* \to G_\tau \equiv \mathrm{Aut}(\mathbb{P}^1, \tau), \qquad [a,b] \to [ax + by, \bar{b}x + \bar{a}y],$$

is an isomorphism. So, G_τ has two topological components, with the automorphism $[x,y] \to [y,x]$ contained in the non-identity component. In general, the map

$$\begin{gathered}
\big((\mathrm{Sym}^d\mathbb{C})^m - \Delta_{\mathbb{R}}\big) \times \mathbb{RP}^{2m-1} \to \mathcal{P}_0(\mathbb{P}^{2m-1}, [d])^{\phi,\eta}, \\
\big([b_{1;1}, \ldots, b_{1;d}], \ldots, [b_{m;1}, \ldots, b_{m;d}], [A_1, \ldots, A_m]\big) \to \\
\Big[A_1 \prod_{r=1}^{d} (x + b_{1;r}y), \bar{A}_1 \prod_{r=1}^{d} (\bar{b}_{1;r}x + y), \ldots, A_m \prod_{r=1}^{d} (x + b_{m;r}y), \bar{A}_m \prod_{r=1}^{d} (\bar{b}_{m;r}x + y)\Big],
\end{gathered}$$

where $\mathbb{R}\mathbb{P}^{2m-1}\equiv(\mathbb{C}^m-0)/\mathbb{R}^*$ and

$$\Delta_{\mathbb{R}}=\big\{([b_{1;1},\ldots,b_{1;d}],\ldots,[b_{m;1},\ldots,b_{m;d}])\in(\mathrm{Sym}^d\mathbb{C})^m:\ S^1\cap\bigcap_{i=1}^{m}\{b_{i;r}:\ r=1,\ldots,d\}\neq\emptyset\in\mathbb{C}\big\},$$

is an isomorphism over the open subset of $\mathcal{P}_0(\mathbb{P}^{2m-1},[d])^{\tau_{2m-1},\tau}$ consisting of maps u such that $u([1,0])$ does not lie in any of the coordinate subspaces of $\mathbb{P}^{2m-1}$. Since the complement of this subspace is of codimension 2, this isomorphism induces orientations on

$$\mathcal{P}_0(\mathbb{P}^{2m-1},[d])^{\tau_{2m-1},\tau}\qquad\text{and}\qquad\mathcal{P}_0(\mathbb{P}^{2m-1},[d])^{\tau_{2m-1},\tau}/G_\tau^0,$$

where $G_\tau^0\subset G_\tau$ is the identity component. On the other hand, the action $\tau_{\mathcal{M}}$ of $[x,y]\to[y,x]$ on $\mathcal{P}_0(\mathbb{P}^{2m-1},[d])^{\tau_{2m-1},\tau}$ lifts over the above orientation-inducing map as

$$\begin{gathered}\big((\mathrm{Sym}^d\mathbb{C})^m-\Delta_{\mathbb{R}}\big)\times\mathbb{R}\mathbb{P}^{2m-1}\to\mathcal{P}_0(\mathbb{P}^{2m-1},[d])^{\tau_{2m-1},\tau},\\ \big([b_{1;1},\ldots,b_{1;d}],\ldots,[b_{m;1},\ldots,b_{m;d}],[A_1,\ldots,A_m]\big)\to\\ \big([b_{1;1}^{-1},\ldots,b_{1;d}^{-1}],\ldots,[b_{m;1}^{-1},\ldots,b_{m;d}^{-1}],[A_1b_{1;1}\ldots b_{1;d},\ldots,A_mb_{m;1}\ldots b_{m;d}]\big).\end{gathered}$$

Since this lift is orientation-preserving, the action of $\tau_{\mathcal{M}}$ on $\mathcal{P}_0(\mathbb{P}^{2m-1},[d])^{\tau_{2m-1},\tau}$ is orientation-preserving and so $\mathcal{M}_l(\mathbb{P}^{2m-1},[d])^{\tau_{2m-1},\tau}$ is orientable for all l. In fact, the resulting orientation on $\mathcal{M}_l(\mathbb{P}^{2m-1},[d])^{\tau_{2m-1},\tau}$ is induced by the canonical spin structure on $T\mathbb{R}\mathbb{P}^{2m-1}$ if m is even and by the canonical relative spin structure on $T\mathbb{R}\mathbb{P}^{2m-1}$ if m is odd; see Remarks A.5 and A.6.

We now show that the conclusion of Lemma 3.3 applies to the moduli spaces $\mathcal{M}_l(\mathbb{P}^{2m-1},[2d])^{\tau_{2m-1},c}$ with the orientations defined in above, provided the left-hand side is suitably oriented. The family of maps

$$\begin{split}[x,y]\to\Big[A_1\prod_{r=1}^{d}(\epsilon a_{1;r}x-y)(x-\epsilon b_{1;r}y),\,\bar{A}_1\prod_{r=1}^{d}(x+\epsilon\bar{a}_{1;r}y)(\epsilon\bar{b}_{1;r}x+y),\ldots,\\ A_m\prod_{r=1}^{d}(\epsilon a_{m;r}x-y)(x-\epsilon b_{m;r}y),\,\bar{A}_m\prod_{r=1}^{d}(x+\epsilon\bar{a}_{m;r}y)(\epsilon\bar{b}_{m;r}x+y)\Big]\end{split}$$

in $\mathcal{M}_l(\mathbb{P}^{2m-1},[2d])^{\eta_{2m-1},\eta}$ with $\epsilon\to0$ converges (before stabilization) to the map

$$u_{\mathrm{top}}\cup u_0\cup u_{\mathrm{bot}}:\mathbb{P}^1_{\mathrm{top}}\cup\mathbb{P}^1_0\cup\mathbb{P}^1_{\mathrm{bot}}\to\mathbb{P}^{2m-1}\qquad\text{with}\quad u_0(\mathbb{P}^1_0)=[(-1)^dA_1,\bar{A}_1,\ldots,(-1)^dA_m,\bar{A}_m],$$

$$u_{\mathrm{top}}([x,y])=\Big[A_1\prod_{r=1}^{d}(a_{1;r}x-y),\,\bar{A}_1\prod_{r=1}^{d}(\bar{b}_{1;r}x+y),\ldots,A_m\prod_{r=1}^{d}(a_{m;r}x-y),\,\bar{A}_m\prod_{r=1}^{d}(\bar{b}_{m;r}x+y)\Big],$$

$$u_{\mathrm{bot}}([x,y])=\Big[A_1\prod_{r=1}^{d}(b_{1;r}y-x),\,\bar{A}_1\prod_{r=1}^{d}(x+\bar{a}_{1;r}y),\ldots,A_m\prod_{r=1}^{d}(b_{m;r}y-x),\,\bar{A}_m\prod_{r=1}^{d}(x+\bar{a}_{m;r}y)\Big].$$

On the other hand, the family of maps

$$\begin{aligned}[x,y]\to\bigg[A_1\prod_{r=1}^{d}(\epsilon a_{1;r}x+y)(x+\epsilon b_{1;r}y),\,\bar{A}_1\prod_{r=1}^{d}(x+\epsilon\bar{a}_{1;r}y)(\epsilon\bar{b}_{1;r}x+y),\ldots,\\ A_m\prod_{r=1}^{d}(\epsilon a_{m;r}x+y)(x+\epsilon b_{m;r}y),\,\bar{A}_m\prod_{r=1}^{d}(x+\epsilon\bar{a}_{m;r}y)(\epsilon\bar{b}_{m;r}x+y)\bigg]\end{aligned}$$

in $\mathcal{M}_l(\mathbb{P}^{2m-1},[2d])^{\tau_{2m-1},\tau}$ with $\epsilon\to 0$ converges (before stabilization) to the map

$$u_{\mathrm{top}}\cup u_0\cup u_{\mathrm{bot}}\colon\mathbb{P}^1_{\mathrm{top}}\cup\mathbb{P}^1_0\cup\mathbb{P}^1_{\mathrm{bot}}\to\mathbb{P}^{2m-1}\qquad\text{with}\quad u_0(\mathbb{P}^1_0)=[A_1,\bar{A}_1,\ldots,A_m,\bar{A}_m],$$

$$u_{\mathrm{top}}([x,y])=\bigg[A_1\prod_{r=1}^{d}(a_{1;r}x+y),\,\bar{A}_1\prod_{r=1}^{d}(\bar{b}_{1;r}x+y),\ldots,A_m\prod_{r=1}^{d}(a_{m;r}x+y),\,\bar{A}_m\prod_{r=1}^{d}(\bar{b}_{m;r}x+y)\bigg],$$

$$u_{\mathrm{bot}}([x,y])=\bigg[A_1\prod_{r=1}^{d}(x+b_{1;r}y),\,\bar{A}_1\prod_{r=1}^{d}(x+\bar{a}_{1;r}y),\ldots,A_m\prod_{r=1}^{d}(x+b_{m;r}y),\,\bar{A}_m\prod_{r=1}^{d}(x+\bar{a}_{m;r}y)\bigg].$$

Thus, there are open subsets

$$\begin{gathered}U\subset\mathcal{M}_{1+l}(\mathbb{P}^{2m-1},[d])\times_{\mathrm{ev}_1}L\times\mathbb{R}^+,\\ W_c\subset\mathcal{M}_l(\mathbb{P}^{2m-1},[2d])^{\tau_{2m-1},c},\qquad\widetilde{W}_c\subset(\mathrm{Sym}^d\mathbb{C})^m\times(\mathrm{Sym}^d\mathbb{C})^m\times\mathbb{RP}^{2m-1},\end{gathered}$$

and a commutative diagram

$$\begin{tikzcd}
& W_\eta \arrow[r] & \widetilde{W}_\eta \arrow[dd] \\
U_c \arrow[ur] \arrow[dr] & & \\
& W_\tau \arrow[r] & \widetilde{W}_\tau
\end{tikzcd}$$

where the diagonal arrows are suitably chosen gluing maps of Lemma 3.3, the horizontal arrows are the inverses of the orientation-inducing maps defined above, and the vertical arrow is the identity in the middle factor and given by

$$\begin{gathered}[a_{1;1},\ldots,a_{1;d},\ldots,a_{m;1},\ldots,a_{m;d}]\to[-a_{1;1},\ldots,-a_{1;d},\ldots,-a_{m;1},\ldots,-a_{m;d}],\\ [A_1,\ldots,A_m]\to[\mathfrak{i}^dA_1,\ldots,\mathfrak{i}^dA_m]\end{gathered}$$

in the first and last factors. Since these two maps are orientation-preserving, both diagonal arrows are orientation-preserving if the left-hand side of the gluing map in Lemma 3.3 is suitably oriented.

A.2. **The canonical orientations.** The first homomorphism $f=(f_1,\ldots,f_n)$ in (1.13) is described by

$$\big\{f_i(\ell,\lambda)\big\}(a_1,\ldots,a_n)=\lambda a_i\ \forall\,(a_0,\ldots,a_n)\!\in\!\ell.$$

With $\mathcal{U}_i\!=\!\{[X_1,\ldots,X_n]\!:\ X_i\!\neq\!0\}$, let

$$\mathbf{z}_i=(z_{i1},\ldots,z_{in}):\mathcal{U}_i\to\mathbb{C}^n,\qquad\text{where}\quad z_{ij}=\frac{X_j}{X_i}\,.$$

The second homomorphism in (1.13) over $\mathcal{U}_i$ is described by

$$(p_1,\ldots,p_n)\to\sum_{j\neq i}\big(p_j(\mathbf{z}_i(\ell))-z_{ij}p_i(\mathbf{z}_i(\ell))\big)\frac{\partial}{\partial z_{ij}}\quad\forall\,p_j\in\mathcal{O}_{\mathbb{P}^{n-1}}(1)|_\ell.$$

It is straightforward to check that this homomorphism is independent of the choice of i and the sequence (1.13) is indeed exact. This short exact sequence gives rise to a natural isomorphism

$$\Lambda^{\mathrm{top}}_{\mathbb{C}}\big(n\mathcal{O}_{\mathbb{P}^{n-1}}(1)\big)\approx\Lambda^{\mathrm{top}}_{\mathbb{C}}\big(\mathbb{P}^{n-1}\!\times\!\mathbb{C}\big)\otimes\Lambda^{\mathrm{top}}_{\mathbb{C}}(T\mathbb{P}^{n-1})\approx K^*_{\mathbb{P}^{n-1}}.\tag{A.1}$$

We define

$$|\phi|=\begin{cases}0, & \text{if } \phi\!=\!\tau_{2m-1};\\ 1, & \text{if } \phi\!=\!\eta_{2m-1};\end{cases}\qquad |c|=\begin{cases}0, & \text{if } c\!=\!\tau;\\ 1, & \text{if } c\!=\!\eta.\end{cases}\tag{A.2}$$

The involution $\phi\!=\!\eta_{2m-1},\tau_{2m-1}$ lifts to an involution Φ on $\mathcal{O}_{\mathbb{P}^{2m-1}}(-1)\!\oplus\!\mathcal{O}_{\mathbb{P}^{2m-1}}(-1)$ by

$$\begin{aligned}\Phi\big(\ell,(x_1,\ldots,x_{2m}),(y_1,\ldots,y_{2m})\big)=\big(\phi(\ell),&(\bar{y}_2,(-1)^{|\phi|}\bar{y}_1,\ldots,\bar{y}_{2m},(-1)^{|\phi|}\bar{y}_{2m-1}),\\ &((-1)^{|\phi|}\bar{x}_2,\bar{x}_1,\ldots,(-1)^{|\phi|}\bar{x}_{2m},\bar{x}_{2m-1})\big).\end{aligned}$$

In turn, this involution induces an involution on the dual bundle $\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\!\oplus\!\mathcal{O}_{\mathbb{P}^{2m-1}}(1)$ by

$$\big\{\Phi(\ell,\alpha_1,\alpha_2)\big\}(\phi(\ell),x,y)=\overline{\big\{(\ell,\alpha_1,\alpha_2)\big\}\big(\Phi(\phi(\ell),x,y)\big)}\tag{A.3}$$

and thus involutions Φ on

$$\mathcal{O}_{\mathbb{P}^{2m-1}}(2)\approx\Lambda^{\mathrm{top}}_{\mathbb{C}}\big(\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\!\oplus\!\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\big),\quad 2m\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\approx m\big(\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\!\oplus\!\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\big),$$

$$\mathcal{O}_{\mathbb{P}^{2m-1}}(2m)\approx\Lambda^{\mathrm{top}}_{\mathbb{C}}\big(2m\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\big)\approx\mathcal{O}_{\mathbb{P}^{2m-1}}(2)^{\otimes m}\tag{A.4}$$

lifting ϕ. The last two lifts commute with the homomorphisms in (1.13) and the isomorphism (A.1), when $n\!=\!2m$ is even.

The isomorphisms (A.1) and (A.4) determine real square root structures on $K_{\mathbb{P}^{4m-1}}$, as needed for orienting the moduli spaces $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi,\eta}$. A spin structure on $\mathbb{R}\mathbb{P}^{4m-1}=\mathrm{Fix}(\tau_{4m-1})$ is determined by a trivialization of $T\mathbb{R}\mathbb{P}^{4m-1}=\mathrm{Fix}(d\tau_{4m-1})$ over any one of the m circles

$$\mathbb{R}\mathbb{P}^1_i=\mathbb{R}\mathbb{P}^1_{\bar{i}}\equiv\big\{[X_1,\ldots,X_{4m}]\!\in\!\mathbb{R}\mathbb{P}^{4m-1}\!:\ X_j\!=\!0\ \forall\,j\!\neq\!i,\bar{i}\big\},\qquad i=1,2,\ldots,2m.$$

Via the real part (the fixed loci of the involutions) of the short exact sequence (1.13), such a trivialization induces a trivialization of

$$\big(4m\mathcal{O}_{\mathbb{P}^{4m-1}}(1)\big)^{\mathbb{R}} \equiv \mathrm{Fix}\big(\Phi\colon 4m\mathcal{O}_{\mathbb{P}^{4m-1}}(1) \to 4m\mathcal{O}_{\mathbb{P}^{4m-1}}(1)\big) \approx 2m\big(2\mathcal{O}_{\mathbb{P}^{4m-1}}(1)\big)^{\mathbb{R}},$$

with the first trivializing section being $f(\cdot,1)$. The homotopy class of the resulting trivialization is independent of the lifts of the $4m-1$ trivializing sections of $T\mathbb{RP}^{4m-1}$ over the homomorphism g in (1.13) and depends only on the homotopy class of the trivialization of $T\mathbb{RP}^{4m-1}$. Furthermore, this induces a bijective correspondence between the homotopy classes of trivializations of the two bundles. On the other hand, any trivialization of $(2\mathcal{O}_{\mathbb{P}^{4m-1}}(1))^{\mathbb{R}}$ over $\mathbb{RP}^1_i$ induces a trivialization of $2(2\mathcal{O}_{\mathbb{P}^{4m-1}}(1))^{\mathbb{R}}$, the homotopy class of which is independent of the choice of the first trivialization. Therefore, there is a canonical homotopy class of trivializations of $(4m\mathcal{O}_{\mathbb{P}^{4m-1}}(1))^{\mathbb{R}}$ over $\mathbb{RP}^1_i$, which in turn determines a homotopy class of trivializations of $T\mathbb{RP}^{4m-1}$ over $\mathbb{RP}^1_i$ and thus a spin structure on $\mathbb{RP}^{4m-1}$ (which is independent of the choice of i). This spin structure determines an orientation on $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1},\tau}$. Since we trivialize the summands $(2\mathcal{O}_{\mathbb{P}^{4m-1}}(1))^{\mathbb{R}}$ in the same way, the orientations on

$$\Lambda^{\mathrm{top}}_{\mathbb{R}}\big(T\mathbb{RP}^{4m-1}\big) = \big(\Lambda^{\mathrm{top}}_{\mathbb{C}}T\mathbb{P}^{4m-1}\big)^{\mathbb{R}} \approx \mathcal{O}_{\mathbb{P}^{4m-1}}(2m)^{\mathbb{R}} \otimes \mathcal{O}_{\mathbb{P}^{4m-1}}(2m)^{\mathbb{R}}$$

and thus on $\mathbb{RP}^{4m-1}$ induced in the two ways described above agree. Therefore, we must flip the canonical orientation of the moduli spaces $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi,\eta}$ when orienting the moduli spaces $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi}$ as in Section 3.

A.3. **Fixed loci.**

For $i=1,2,\ldots,2m$, we define

$$\bar{i} = \begin{cases} i+1, & \text{if } 2\not| i; \\ i-1, & \text{if } 2|i. \end{cases}$$

The m-torus $\mathbb{T}$ acts on $\mathbb{P}^{2m-1}$ by

$$(\zeta_1,\ldots,\zeta_m)\cdot[z_1,z_2,\ldots,z_{2m-1},z_{2m}] = [\zeta_1 z_1,\zeta_1^{-1}z_2,\ldots,\zeta_m z_{2m-1},\zeta_m^{-1}z_{2m}].$$

This action commutes with the involutions $\phi=\tau_{2m-1},\eta_{2m-1}$ and has $2m$ fixed points,

$$p_1 = [1,0,\ldots,0], \qquad \ldots \qquad p_{2m} = [0,\ldots,1].$$

We note that $\phi(p_i)=p_{\bar{i}}$. By composition on the left, $\mathbb{T}$ also acts on $\mathcal{M}_l(\mathbb{P}^{2m-1},[d])^{\phi,c}$, where $c=\tau,\eta$.

Lemma A.1 ([16, Lemma 3.1])**.** *The irreducible $\mathbb{T}$-fixed curves in $\mathbb{P}^{2m-1}$ are the lines L_{ij} connecting the points p_i and p_j with $i\neq j$. Moreover, the irreducible ϕ- and $\mathbb{T}$-fixed curves in $\mathbb{P}^{2m-1}$ are the lines $L_{i\bar{i}}$.*

Let $\lambda_i \in H^*_{\mathbb{T}}$ be the equivariant first Chern class of $\mathcal{O}_{\mathbb{P}^{2m-1}}|_{p_i}$. Thus,

$$\lambda_{\bar{i}} = -\lambda_i, \qquad H^*_{\mathbb{T}} = \mathbb{Q}[\lambda_1,\lambda_3,\ldots,\lambda_{2m-1}].$$

Let $[f,(z_k,c(z_k))_k]$ be an element of $\mathcal{M}_l(\mathbb{P}^{2m-1},[d])^{\phi,c}$ fixed by the $\mathbb{T}$-action. Since there are no $\mathbb{T}$-fixed points in $\mathbb{P}^{2m-1}$ that are also fixed by ϕ, the domain Σ of f contains a



central component Σ_0, while the remaining irreducible components come in conjugate pairs. Furthermore, $f_0 \equiv f|_{\Sigma_0}$ is a cover of some line $L_{i\bar{i}}$ of some degree $d_0 \in \mathbb{Z}^+$ which is branched only over p_i and $p_{\bar{i}}$. Every nodal and marked point of Σ and branched point of f is mapped to a fixed point p_j. If $d_0 < d$ or $l > 1$, the complement of Σ_0 in Σ consists of two nodal curves Σ' and Σ'', each with $l+1$ marked points $(x_k)_{k=0}^{l}$ so that x_0 corresponds to the node shared with Σ_0 and each of the remaining points is decorated by a sign s_k, $+$ or $-$, depending on whether it is the first or the second point in the pair $(z_k, c(z_k))$.

Similarly to [10, Section 27.3], every fixed locus of such maps can be modeled on a labeled tree, Γ, symmetric about the mid-point of a distinguished edge e_0, which corresponds to the central component Σ_0 of the $\mathbb{T}$-fixed maps in the locus. Every edge e of Γ is labeled by some $d_e \in \mathbb{Z}^+$, indicating the degree of the corresponding map; these labels are preserved by the reflection symmetry of Γ. Every vertex v is labeled by some $j_v = 1, 2, \ldots, 2m$ in such a way that the reflection symmetry takes a vertex labeled j to a vertex labeled $\bar{j}$. The graph Γ also contains open edges which correspond to the marked points of the domain Σ; we denote by $v(k)$ the vertex to which the k-th marked point is attached. Figure A.3(a) shows one such graph describing a $\mathbb{T}$-locus in $\overline{\mathcal{M}}_7(\mathbb{P}^{2m-1}, [7])^{\phi,c}$. Removing e_0 from Γ, we get a disconnected graph $\Gamma' \sqcup \bar{\Gamma}'$, where $\bar{\Gamma}'$ is the graph obtained from Γ' by replacing each vertex label j by $\bar{j}$. Choose one of the connected subgraphs, e.g. Γ' , and add the corresponding half-edge in place of the central edge; see Figure A.3(b). We denote the total half graph by Γ_{half}. All calculations below are based on this half-graph; it is straightforward to check that the result is independent of which half we choose.

For each vertex v in Γ_{half}, let $\overline{\mathcal{M}}_v = \overline{\mathcal{M}}_{0,\text{val}(v)}$, where $\text{val}(v)$ is the valence of v, i.e. the number of edges and open edges in Γ leaving v, and $\overline{\mathcal{M}}_{0,\text{val}(v)}$ denotes a point if $\text{val}(v) = 1, 2$. Let

$$\overline{\mathcal{M}}_{\Gamma_{\text{half}}} = \prod_{v} \overline{\mathcal{M}}_v\,, \qquad D_{\Gamma_{\text{half}}} = d_0 \cdot \prod_{e} d_e,$$

where the products are taken over the vertices v and edges e in Γ_{half}.

A.4. **Normal bundles.** For every flag $F = (v, e)$, let $j_F = j_v$. For every element $[f, (z_k, c(z_k))_k]$ in the fixed locus corresponding to Γ, there is an exact sequence

$$0 \to \text{Aut}\big(\Sigma, (z_k, c(z_k))_k\big)_{\mathbb{R}} \to \text{Def}(f)_{\mathbb{R}} \to \text{Def}\big(f, (z_k, c(z_k))_k\big)_{\mathbb{R}} \to \text{Def}(\Sigma, (z_k, c(z_k))_k)_{\mathbb{R}} \to 0,$$

where Σ is the domain of f. Thus,

$$e(N_\Gamma)=e\big(\mathrm{Def}\big(f,(z_k,c(z_k))_k\big)^{\mathrm{mov}}_{\mathbb{R}}\big)=\frac{e(\mathrm{Def}(f)^{\mathrm{mov}}_{\mathbb{R}})\,e(\mathrm{Def}(\Sigma,(z_k,c(z_k))_k)^{\mathrm{mov}}_{\mathbb{R}})}{e(\mathrm{Aut}(\Sigma,(z_k,c(z_k))_k)^{\mathrm{mov}}_{\mathbb{R}})},\tag{A.5}$$

where "mov" means the moving part (the part with the nonzero $\mathbb{T}$-weights) and $e(\cdot)$ denotes the equivariant Euler class. Following [10, Section 27.4], we now determine the three terms appearing on the right-hand side of (A.5).

For each edge e of Γ_{half}, $\mathrm{Aut}(\Sigma,(z_k,c(z_k))_k)_{\mathbb{R}}$ contains a $\mathbb{T}$-fixed one-dimensional complex subspace of infinitesimal automorphisms of the corresponding non-contracted component Σ_e which fix the two branch points of $f_e\equiv f|_{\Sigma_e}$; this subspace cancels with a similar piece in $\mathrm{Def}(f_e)_{\mathbb{R}}$. The space $\mathrm{Aut}(\Sigma,(z_k,c(z_k))_k)_{\mathbb{R}}$ also contains a $\mathbb{T}$-fixed one-dimensional real subspace of infinitesimal automorphisms of the central component Σ_0; this subspace cancels with a similar piece in $\mathrm{Def}(f_0)_{\mathbb{R}}$, up to sign taken into account by Lemma A.2. The remaining automorphisms, none of which is $\mathbb{T}$-fixed, correspond to the vertices v in Γ_{half} of valence 1; they describe the infinitesimal automorphisms moving the branch point x_v of f_e, where e is the unique edge containing v, that lies over j_v. Thus, similarly to [10, Section 27.4],

$$\begin{gathered}e\big(\mathrm{Aut}(\Sigma,(z_k,c(z_k))_k)^{\mathrm{mov}}_{\mathbb{R}}\big)=\prod_{\substack{v\in e\\ \mathrm{val}(v)=1}}e(T_{x_v}\Sigma_e)=\prod_{\substack{v\in e\\ \mathrm{val}(v)=1}}w_{(v,e)}\,,\\ \text{where}\qquad w_{(v,\{v,v'\})}=\frac{\lambda_{j_v}-\lambda_{j_{v'}}}{d_{\{v,v'\}}}\,.\end{gathered}\tag{A.6}$$

A deformation of a contracted component of the domain (as a marked curve) is $\mathbb{T}$-fixed. The moving deformations come from smoothing (conjugate pairs) of nodes of Σ. For each node x of Σ corresponding to Γ_{half}, $\mathrm{Def}(\Sigma,(z_k,c(z_k))_k)^{\mathrm{mov}}_{\mathbb{R}}$ contains the complex one-dimensional space isomorphic to the tensor product of the tangent spaces of the two components of Σ sharing x. There are two possibilities. Each $v\in\Gamma_{\mathrm{half}}$ shared by two edges contributes $w_{F_1}+w_{F_2}$, where F_1 and F_2 are the two flags containing v. Each flag $F=(v,e)$ with $v\in\Gamma_{\mathrm{half}}$ and $\mathrm{val}(v)\geq 3$ contributes $w_F-\psi_F$, where $\psi_F\in H^2(\overline{\mathcal{M}}_v)$ is the first Chern class of the universal cotangent bundle on $\overline{\mathcal{M}}_v$ corresponding to the marked point determined by F on the contracted curve determined by the vertex v. Thus,

$$e\big(\mathrm{Def}(\Sigma,(z_k,c(z_k))_k)^{\mathrm{mov}}_{\mathbb{R}}\big)=\prod_{\substack{\mathrm{val}(v)=2\\ v\in e_1,e_2\\ e_1\neq e_2}}\big(w_{(v,e_1)}+w_{(v,e_2)}\big)\cdot\prod_{\mathrm{val}(v)\geq 3}\prod_{v\in e}\big(w_{(v,e)}-\psi_{(v,e)}\big).\tag{A.7}$$

Finally, there is an exact sequence

$$\begin{aligned}0\to\mathrm{Def}(f)_{\mathbb{R}}\to H^0(\Sigma_{e_0},f_0^*T\mathbb{P}^{2m-1})_{\mathbb{R}}\oplus\bigoplus_{e\neq e_0}H^0(\Sigma_e,f_e^*T\mathbb{P}^{2m-1})\oplus\bigoplus_{v}T_{p_{j_v}}\mathbb{P}^{2m-1}\\ \to\bigoplus_{F}T_{p_{j_F}}\mathbb{P}^{2m-1}\to 0,\end{aligned}$$

where the direct sums are taken over the vertices v, edges e, and flags F in Γ_{half}. Thus,

$$e\big(\mathrm{Def}(f)^{\mathrm{mov}}_{\mathbb{R}}\big)=\frac{e(H^0(\Sigma_{e_0},f_0^*T\mathbb{P}^{2m-1})^{\mathrm{mov}}_{\mathbb{R}})\prod\limits_{e\neq e_0}e(H^0(\Sigma_e,f_e^*T\mathbb{P}^{2m-1})^{\mathrm{mov}})\cdot\prod\limits_{v}\prod\limits_{j\neq j_v}(\lambda_{j_v}-\lambda_j)}{\prod\limits_{F}\prod\limits_{j\neq j_F}(\lambda_{j_F}-\lambda_j)}\,. \tag{A.8}$$

The contribution of $e\neq e_0$ is standard and given by

$$e\big(H^0(\Sigma_e,f_e^*T\mathbb{P}^{2m-1})^{\mathrm{mov}}\big)=(-1)^{d_e}\frac{d_e!^2}{d_e^{2d_e}}(\lambda_{j_1}-\lambda_{j_2})^{2d_e}\prod_{r=0}^{d_e}\prod_{k\neq j_1,j_2}\bigg(\frac{r\lambda_{j_1}+(d_e-r)\lambda_{j_2}}{d_e}-\lambda_k\bigg)\,, \tag{A.9}$$

where j_1 and j_2 are the two vertex labels of the edge e; see [10, Section 27.4]. The contribution of the half-edge e_0 is described by the next lemma, which is proved in Section A.6.

Lemma A.2. *Let* $\phi=\tau_{4m-1},\eta_{4m-1}$, $c=\tau,\eta$, *and* $f_0\colon(\mathbb{P}^1,[1,0],[0,1])\to(\mathbb{P}^{4m-1},p_i,p_{\bar{i}})$ *be the degree* d_0 *cover of a line* $L_{i\bar{i}}$ *branched over only* p_i *and* $p_{\bar{i}}$ *and intertwining the involutions* c *and* ϕ. *With respect to the canonical orientation of the moduli space* $\overline{\mathcal{M}}_0(\mathbb{P}^{4m-1},[d_0])^{\phi,c}$,

$$e\big(H^0(\Sigma_{e_0},f_0^*T\mathbb{P}^{4m-1})^{\mathrm{mov}}_{\mathbb{R}}\big)=d_0!\bigg(\frac{2\lambda_i}{d_0}\bigg)^{d_0}\prod_{\substack{1\le j\le 4m\\ j\neq i,\,2|(i+j)}}\prod_{r=0}^{d_0}\bigg(\frac{d_0-2r}{d_0}\lambda_i-\lambda_j\bigg). \tag{A.10}$$

A.5. **Applications.** By the classical localization theorem of [1],

$$N_d^{\phi}(t_1,\ldots,t_l)=\sum_{\Gamma\rhd\tau}\frac{1}{D_{\Gamma_{\mathrm{half}}}}\int_{\mathcal{M}_{\Gamma_{\mathrm{half}}}}\frac{\prod\limits_{k=1}^{l}s_k^{t_k+1}\lambda_{j_{v(k)}}^{t_k}}{e(N_\Gamma)}-\sum_{\Gamma\rhd\eta}\frac{1}{D_{\Gamma_{\mathrm{half}}}}\int_{\mathcal{M}_{\Gamma_{\mathrm{half}}}}\frac{\prod\limits_{k=1}^{l}s_k^{t_k+1}\lambda_{j_{v(k)}}^{t_k}}{e(N_\Gamma)}\,, \tag{A.11}$$

where the first and second sums are taken over the graphs Γ corresponding to the fixed loci in $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi,\tau}$ and $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi,\eta}$, respectively. The negative sign arises due to the fact we flip the orientation of $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi,\eta}$ when gluing it to $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi,\tau}$. Along with (A.5)-(A.10), (A.11) provides an explicit way of computing the numbers (1.14).

Proof of Theorem 1.8. Suppose t_k is even for some k. Given any graph Γ corresponding to a fixed locus in $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\phi,c}$, let Γ' be the graph obtained from Γ by changing the sign of the k-th marked point. By (A.11), the contribution of Γ' to $N_d^{\phi}(t_1,\ldots,t_l)$ is the negative of the contribution of Γ. Thus, $N_d^{\phi}(t_1,\ldots,t_l)$ vanishes if some t_k is even.

If d is even, the graphs Γ corresponding to the fixed loci in $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1},\tau}$ and $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1},\eta}$ are the same. By (A.11), this implies that $N_d^{\tau_{4m-1}}(t_1,\ldots,t_l)$ vanishes if d is even.

If d is odd, the graphs Γ corresponding to the fixed loci in $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1},\tau}$ and $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\eta_{4m-1},\eta}$ are the same. Along with (1.16) and (A.110, this implies (1.15). $\square$

If $d, t_1, \ldots, t_l$ are odd, (A.11) gives

$$N_d^{\tau_{4m-1}}(t_1,\ldots,t_l) = 2^{l-1}\sum_{\Gamma_{\mathrm{half}}} \frac{1}{D_{\Gamma_{\mathrm{half}}}} \int_{\mathcal{M}_{\Gamma_{\mathrm{half}}}} \frac{\prod\limits_{k=1}^{l} \lambda_{j_{v(k)}}^{t_k}}{e(N_\Gamma)} = -N_d^{\eta_{4m-1}}(t_1,\ldots,t_l), \tag{A.12}$$

where the sum is taken over all half-graphs Γ_{half} corresponding to $\overline{\mathcal{M}}_l(\mathbb{P}^{4m-1},[d])^{\tau_{4m-1},\tau}$ containing marked points with the $+$ sign only.

Example A.3 (d=1)**.** If $d=1$, $t_1,\ldots,t_l \in \mathbb{Z}^+$ are odd, and $t_1+\ldots+t_l = 4m-2+l$, (A.12) gives

$$N_1^{\tau_{4m-1}}(t_1,\ldots,t_l) = -N_1^{\eta_{4m-1}}(t_1,\ldots,t_l) = 2^l \sum_{i=1}^{2m} \frac{\lambda_i^{t_1+\ldots+t_l}}{2\lambda_i \prod\limits_{\substack{1\le j\le 2m\\ j\neq i}} (\lambda_j^2-\lambda_i^2)} (2\lambda_i)^{-(l-1)}.$$

Simplifying and using the residue theorem on S^2, we obtain

$$N_1^{\tau_{4m-1}}(t_1,\ldots,t_l) = -N_1^{\eta_{4m-1}}(t_1,\ldots,t_l) = -\sum_{i=1}^{2m} \operatorname*{Res}_{z=\lambda_i^2} \frac{z^{2m-1}dz}{\prod\limits_{j=1}^{2m}(z-\lambda_j^2)} = \operatorname*{Res}_{z=\infty} \frac{z^{2m-1}dz}{\prod\limits_{j=1}^{2m}(z-\lambda_j^2)} = -1.$$

Example A.4 (m=1,d=3)**.** There are five distinct types of half-graphs that contribute to the numbers $N_3^{\phi}(3,3,3)$ with $\phi=\tau_3,\eta_3$. In this case, we obtain

$$\begin{aligned} N_3^{\tau_3}(3,3,3) = 4\Bigg[&\left(\frac{3}{2}\right)^5 \left(\frac{\lambda_1^4}{(\lambda_1^2-\lambda_3^2)(\lambda_1^2-9\lambda_3^2)} + \frac{\lambda_3^4}{(\lambda_3^2-\lambda_1^2)(\lambda_3^2-9\lambda_1^2)}\right) \\ &- \frac{1}{32}\frac{\lambda_1^4+\lambda_3^4}{(\lambda_1^2-\lambda_3^2)^2} + \frac{1}{(\lambda_1^2-\lambda_3^2)^4}\left(\frac{\lambda_3^8(3\lambda_1^2+5\lambda_3^2)}{9\lambda_1^2-\lambda_3^2} + \frac{\lambda_1^8(3\lambda_3^2+5\lambda_1^2)}{9\lambda_3^2-\lambda_1^2}\right) \\ &- \frac{3}{16}\frac{\lambda_1^4+\lambda_3^4}{(\lambda_1^2-\lambda_3^2)^2} - 3\frac{\lambda_1^2\lambda_3^2(\lambda_1^4+\lambda_3^4)}{(\lambda_1^2-\lambda_3^2)^4} - \frac{3}{8}\frac{\lambda_1^4+\lambda_3^4}{(\lambda_1^2-\lambda_3^2)^2} + \frac{3}{4}\frac{\lambda_1^2\lambda_3^2(3\lambda_1^4+2\lambda_1^2\lambda_3^2+3\lambda_3^4)}{(\lambda_1^2-\lambda_3^2)^4} \\ &- \frac{1}{4}\frac{\lambda_1^4+\lambda_3^4}{(\lambda_1^2-\lambda_3^2)^2} - \frac{1}{2}\frac{\lambda_1^6(3\lambda_1^2-\lambda_3^2)+\lambda_3^6(3\lambda_3^2-\lambda_1^2)}{(\lambda_1^2-\lambda_3^2)^4}\Bigg] = 1. \end{aligned}$$

A.6. **Proof of Lemma A.2.** The restriction of

$$\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\otimes\mathcal{O}_{\mathbb{P}^{2m-1}}(1) \approx \Lambda_{\mathbb{C}}^{\mathrm{top}}\big(\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\oplus\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\big)$$

to $L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\}$ is trivialized by the homomorphism

$$(\ell,\alpha) \to \big(\ell, \mathfrak{i} z_{i\bar{i}}^{-1}\alpha(\mathbf{z}_i(\ell)\otimes\mathbf{z}_i(\ell))\big).$$

Via this trivialization, the involution Φ on $\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\otimes\mathcal{O}_{\mathbb{P}^{2m-1}}(1)$ induced by (A.3), i.e.

$$\begin{split}&\big\{\Phi(\ell,\alpha)\big\}\big(\phi(\ell),(x_1,\ldots,x_{2m})\otimes(y_1,\ldots,y_{2m})\big)\\ &=-\overline{\{\alpha_1\otimes\alpha_2\}\big((\bar{y}_2,(-1)^{|\phi|}\bar{y}_1,\ldots,\bar{y}_{2m},(-1)^{|\phi|}\bar{y}_{2m-1})\otimes((-1)^{|\phi|}\bar{x}_2,\bar{x}_1,\ldots,(-1)^{|\phi|}\bar{x}_{2m},\bar{x}_{2m-1})\big)},\end{split}$$

corresponds to the standard involution on $(L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\})\times\mathbb{C}$ lifting the restriction of ϕ. Thus, the trivialization

$$\mathcal{O}_{\mathbb{P}^{2m-1}}(2m)\big|_{L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\}}\to(L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\})\times\mathbb{C},\qquad(\ell,\alpha)\to\big(\ell,\mathfrak{i}^m z_{i\bar{i}}^{-m}\alpha(\mathbf{z}_i(\ell)^{\otimes 2m})\big),\tag{A.13}$$

is an admissible trivialization of $(\mathcal{O}_{\mathbb{P}^{2m-1}}(2m),\Phi)$ induced by a trivialization of its real square root via the canonical isomorphism

$$\mathcal{O}_{\mathbb{P}^{2m-1}}(2m)\approx\mathcal{O}_{\mathbb{P}^{2m-1}}(m)\otimes\mathcal{O}_{\mathbb{P}^{2m-1}}(m)$$

if m is even.

In the case $\phi=\tau_{2m-1}$,

$$\begin{split}\big(2\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\big)^{\mathbb{R}}=\big\{(\alpha_1,\alpha_2)\in 2\mathcal{O}_{\mathbb{P}^{2m-1}}(1)|_\ell\colon\ \ell\in\mathbb{RP}^{2m-1},&\\ \alpha_2(x_1,\ldots,x_{2m})=\overline{\alpha_1(\bar{x}_2,\bar{x}_1,\ldots,\bar{x}_{2m-1},\bar{x}_{2m})}\ \forall\,(x_1,\ldots,x_{2m})\in\ell&\big\}.\end{split}$$

Thus, we can trivialize $(2\mathcal{O}_{\mathbb{P}^{2m-1}}(1))^{\mathbb{R}}$ over $\mathbb{RP}^1_i$ by

$$(\ell,\alpha_1,\alpha_2)\to\big(\ell,\alpha_1(\mathbf{z}_i(\ell))\big)\in\mathbb{RP}^1_i\times\mathbb{C}.\tag{A.14}$$

Thus, by the discussion at the end of Section A.2, the trivialization

$$\Psi\colon\big(2m\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\big)^{\mathbb{R}}\to\mathbb{RP}^1_i\times\mathbb{C}^{\{j\colon 2|(j-i)\}},\qquad\Psi(\ell,\alpha_1,\ldots,\alpha_{2m})_j=\alpha_j(\mathbf{z}_i(\ell)),\tag{A.15}$$

determines the canonical spin structure on $\mathbb{RP}^{2m-1}$ if m is even.

The composition of the lift $\tilde{\phi}$ of ϕ to the standard conjugation on $L_{i\bar{i}}\times\mathbb{C}^{2m-1}$ with the conjugation Φ_i induced by the trivialization of $T\mathbb{P}^{2m-1}$ along $L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\}$ by the standard coordinate vector fields as in Section A.2 is given by

$$\tilde{\phi}\circ\Phi_i\colon\mathbb{C}^*\to\mathrm{GL}(2m-1,\mathbb{C}),\quad\big(\tilde{\phi}\circ\Phi_i(z_{i\bar{i}})\big)_{j_1j_2}=\begin{cases}(-1)^{|\phi|+1}z_{i\bar{i}}^{-2}\,, & \text{if } j_1,j_2=\bar{i};\\ (-1)^{|\phi|(i+j_1)}z_{i\bar{i}}^{-1}\,, & \text{if } j_1=\bar{j}_2\neq i,\bar{i};\\ 0, & \text{otherwise.}\end{cases}\tag{A.16}$$

Thus, the composition of this trivialization with (id,A), where

$$A_i\colon L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\}\to\mathrm{GL}(2m-1,\mathbb{C}),\ \big(A_i(z_{i\bar{i}})\big)_{j_1j_2}=\begin{cases}\mathfrak{i}az_{i\bar{i}}^{-1}\,, & \text{if } j_1,j_2=\bar{i};\\ (-\mathfrak{i})^{j_1+j_2+1}, & \text{if } j_1\in\{j_2,\bar{j}_2\},2|(j_2-i);\\ \mathfrak{i}^{j_1+j_2}z_{i\bar{i}}^{-1}, & \text{if } j_1\in\{j_2,\bar{j}_2\},2|(j_2-\bar{i});\\ 0, & \text{otherwise;}\end{cases}\tag{A.17}$$

corresponds to the standard involution on $(L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\})\times\mathbb{C}^{2m-1}$ lifting ϕ, whenever $a\in\mathbb{R}^*$.

We order the standard coordinate vector fields along $L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\}$ so that $\frac{\partial}{\partial z_{i\bar{i}}}$ is listed first, followed by the pairs consisting of $\frac{\partial}{\partial z_{ij}}$ and $\frac{\partial}{\partial z_{i\bar{j}}}$ with $j\neq i$ and $2|(j-i)$. The corresponding element of $\Lambda^{\mathrm{top}}_{\mathbb{C}}(T\mathbb{P}^{2m-1})$ is then mapped to

$$\det\big(A_i(z_{i\bar{i}})\big)=\mathfrak{i}^m 2^{m-1} a z_{i\bar{i}}^{-m}$$

under the trivialization of $\Lambda^{\mathrm{top}}_{\mathbb{C}}(T\mathbb{P}^{2m-1})$ over $L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\}$ induced by the composite trivialization of $T\mathbb{P}^{2m-1}$. On the other hand, the image of this $(2m-1)$-tensor under the canonical isomorphism (A.1) followed by the trivialization (A.13) is $(-1)^{m\bar{i}}\mathfrak{i}^m z_{i\bar{i}}^{-m}$. Thus, if m is even, the two trivializations of $(K^*_{\mathbb{P}^{n-1}},\Phi)$ over $L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\}$ are homotopic in the sense of Definition 2.1 if and only if $a>0$.

In the case $\phi=\tau_{2m-1}$, the above composite trivialization restricts to a trivialization of $T\mathbb{RP}^{2m-1}$ over $\mathbb{RP}^1_i$. The first trivializing section is $-\mathfrak{i}a^{-1}z_{i\bar{i}}\frac{\partial}{\partial z_{i\bar{i}}}$, followed by

$$\frac{1}{2}\bigg(-\mathfrak{i}\frac{\partial}{\partial z_{ij}}+\mathfrak{i}z_{i\bar{i}}\frac{\partial}{\partial z_{i\bar{j}}}\bigg)\qquad\text{and}\qquad\frac{1}{2}\bigg(\frac{\partial}{\partial z_{ij}}+z_{i\bar{i}}\frac{\partial}{\partial z_{i\bar{j}}}\bigg)\tag{A.18}$$

with $j\neq i$ and $2|(j-i)$. Lifting these sections over the homomorphism g in the real part of the short exact sequence (1.13) and combining with the image of f, we obtain a trivialization of $(2m\mathcal{O}_{\mathbb{P}^{2m-1}})^{\mathbb{R}}$. The composition of this trivialization with the trivialization (A.15) sends the two standard real basis elements of each factor of $\mathbb{C}$ to either 1 and $\mathfrak{i}/2a$ or $-\mathfrak{i}/2$ and $1/2$ in some factor of $\mathbb{C}$. Therefore, if m is even, the trivialization of $(2m\mathcal{O}_{\mathbb{P}^{2m-1}})^{\mathbb{R}}$ over $\mathbb{RP}^1_i$ just discussed differs from a canonical one by a constant matrix-valued function which is orientation-preserving if and only if $a>0$. Thus, the trivialization of $T\mathbb{RP}^{2m-1}$ over $\mathbb{RP}^1_i$ induced by the above composite trivialization corresponds to the canonical spin structure on $\mathbb{RP}^{2m-1}$ if $a>0$ and m is even.

By (A.16), the components of a section $s\in H^0(\Sigma_0,f_0^*T\mathbb{P}^{2m-1})_{\mathbb{R}}$ with respect to the trivialization of $f_0^*T\mathbb{P}^n$ over $\mathbb{P}^1\backslash\{0,\infty\}$ induced by the coordinate tangent vectors along $L_{i\bar{i}}\backslash\{p_i,p_{\bar{i}}\}$ satisfy

$$\begin{aligned}
s_{\bar{i}}(z)&=z^{d_0}\sum_{r=-d_0}^{d_0}s_{\bar{i};r}z^r, &\quad s_{\bar{i};r}&=(-1)^{1+|c|r}\overline{s_{\bar{i};-r}};\\
s_j(z)&=\sum_{r=0}^{d_0}s_{j;r}z^r, &\quad s_{j;r}&=(-1)^{|\phi|(i+j)+|c|r}\overline{s_{\bar{j};d_0-r}} \qquad \forall\, j\neq i,\bar{i}.
\end{aligned}$$

Therefore, the complex coefficients $s_{\bar{i};-r}$, with $r=1,\ldots,d_0$, and $s_{j;-r}$, with $r=0,1,\ldots,d_0$, $j\neq i$, and $2|(j-i)$, and the real coefficient $\mathfrak{i}s_{\bar{i};0}$ give coordinates on $H^0(\Sigma_0,f_0^*T\mathbb{P}^{2m-1})_{\mathbb{R}}$. With

A_i as in (A.17), the components of $A_i s$ thus satisfy

$$(A.19)\qquad (A_i s)_j(z)=\sum_{r=-d_0}^{d_0} b_{j;r}z^r,\quad b_{\bar{i};r}=(-1)^{|c|r}\overline{b_{\bar{i};-r}},\qquad \text{with}\quad b_{\bar{i};r}=\mathfrak{i}as_{\bar{i};r},$$

$$b_{j;0}=2\begin{cases}-\mathrm{Im}(s_{j;0}), & \text{if } j\neq i,\ 2|(j-i);\\ \mathrm{Re}(s_{\bar{j}};0), & \text{if } \bar{j}\neq i,\ 2|(\bar{j}-i);\end{cases}\qquad (b_{j;r},b_{j;-r})=\begin{cases}\mathfrak{i}(s_{j;r},-s_{\bar{j};d_0-r}), & \text{if } r\neq 0,\ j\neq i,\ 2|(j-i);\\ (s_{\bar{j};r},s_{j;d_0-r}), & \text{if } r\neq 0, \bar{j}\neq i,\ 2|(\bar{j}-i).\end{cases}$$

Under the collapsing procedures of Lemma 2.3 and [7, Proposition 8.1.4], the parts of $A_i s$ involving positive and negative powers of z correspond to holomorphic sections on Σ_{top} and Σ_{bot}, respectively. The weights of the $\mathbb{T}$-actions on the space of sections $\{s_{\bar{i};r}z^r\}$ with $1\le r\le d_0$ and $\{s_{j;r}z^r\}$ with $j\neq i,\bar{i}$ and $1\le r\le d_0$ are

$$2\lambda_i\left(1-\frac{r}{d_0}\right)\qquad\text{and}\qquad \frac{d_0-2r}{d_0}\lambda_i-\lambda_j\,,$$

respectively. Since we use the complex orientation of sections on Σ_{top}, these parts contribute

$$d_0!\left(\frac{2\lambda_i}{d_0}\right)^{d_0}\prod_{\substack{1\le j\le 2m\\ j\neq i,\,2|(i+j)}}\prod_{r=1}^{d_0}\left(\frac{d_0-2r}{d_0}\lambda_i-\lambda_j\right)$$

to $e(H^0(\Sigma_{e_0},f_0^*T\mathbb{P}^{4m-1})_{\mathbb{R}}^{\mathrm{mov}})$. The parts of $A_i s$ constant in z correspond to holomorphic sections on Σ_0 commuting with the involution and constitute the direct sum of the trivial representation of $\mathbb{T}$ on the space of sections $\{b_{\bar{i};0}z^0:\ b_{\bar{i};0}\in\mathbb{R}\}$ and of the two-dimensional representations of weight λ_i–λ_j with $j\neq i$ and $2|(j$–$i)$. By the above, the resulting orientation of $H^0(\Sigma_0,f_0^*T\mathbb{P}^{2m-1})_{\mathbb{R}}$ agrees with the one induced by the chosen square root (for $c=\eta$) or the chosen spin structure (for $c=\tau$) if and only if $a>0$.

The characteristic vector field s for the action of $S^1\subset G_c$ which fixes the point $z=0$ in $\mathbb{C}\subset\mathbb{P}^1$ is given by

$$(A.20)\qquad (A_i s)_{\bar{i}}=\mathfrak{i}a\,z^{-d_0}\frac{d}{d\theta}\big(e^{-\mathfrak{i}\theta}z\big)^{d_0}\Big|_{\theta=0}=ad_0,\qquad (A_i s)_j=0\ \forall\, j\neq i,\bar{i}.$$

Since $ad_0>0$ whenever $a>0$, this concludes the proof of Lemma A.2.

Remark A.5. Let $c=\tau,\eta$, $\phi=\tau_{2m-1},\eta_{2m-1}$, and $d\in\mathbb{Z}^+$ be such that $2|(d|c|-|\phi|)$, as in Section A.1. If i is odd, the orientation on $H^0(\Sigma_{e_0},f_0^*T\mathbb{P}^{2m-1})_{\mathbb{R}}^{\mathrm{mov}}$ provided by the above composite trivialization involving the matrix A_i with $a>0$ agrees with the orientation of the tangent space of $\mathcal{P}_0(\mathbb{P}^{2m-1},[d_0])^{\phi,c}$ at f_0 defined in Section A.1. In particular, the deformations of the coefficients of z^r in

$$A_j\prod_{r=1}^{d}(1\mp b_{j;r}z)\Big/A_i\prod_{r=1}^{d}(1\mp b_{i;r}z)$$

for $j\neq i$ and $2|(j-i)$ correspond to the coefficients $b_{j;r}$ in (A.19), up to a complex multiple. The deformations of the coefficients of z^{d+r} in

$$\bar{A}_i\prod_{r=1}^{d}(\bar{b}_{i;r}+z)\Big/A_i\prod_{r=1}^{d}(1\mp b_{i;r}z)$$

with $r>0$ correspond to the coefficients $b_{\bar{i};r}$ in (A.19). Finally, the characteristic vector field corresponding to the S^1 part of the variation of A_i is $-2\mathfrak{i}z^d\frac{\partial}{\partial z_{i\bar{i}}}$, which corresponds to $2a>0$ in $\mathbb{R}$. Thus, if i is even, the orientation on $H^0(\Sigma_{e_0},f_0^*T\mathbb{P}^{2m-1})^{\mathrm{mov}}_{\mathbb{R}}$ provided by the above composite trivialization involving the matrix A_i with $a>0$ agrees with the orientation of the tangent space of $\mathcal{P}_0(\mathbb{P}^{2m-1},[d_0])^{\phi,c}$ at f_0 defined by the map

$$\begin{aligned}&\big([a_{1;1},\ldots,a_{1;d}],\ldots,[a_{m;1},\ldots,a_{m;d}],[B_1,\ldots,B_m]\big)\\&\qquad\to\bigg[\bar{B}_1\prod_{r=1}^{d}(\bar{a}_{1;r}x-y),B_1\prod_{r=1}^{d}(x+a_{1;r}y),\ldots,\bar{B}_m\prod_{r=1}^{d}(\bar{a}_{m;r}x-y),B_m\prod_{r=1}^{d}(x+a_{m;r}y)\bigg].\end{aligned}$$

The two orientations on $\mathcal{P}_0(\mathbb{P}^{2m-1},[d_0])^{\phi,c}$ differ by $(-1)^{(d_0+1)m}$. Thus, if $\mathcal{P}_0(\mathbb{P}^{2m-1},[d_0])^{\phi,c}$ is oriented as in Section A.1, (A.10) extends to $\mathbb{P}^{2m-1}$ in place of $\mathbb{P}^{4m-1}$ by multiplying its right-hand side by $(-1)^{(d_0+1)m\bar{i}}$. The resulting contribution to $N_d^{\phi}(t_1,\ldots,t_l)$ is still independent of the choice of Γ_{half}. By the results of Section A.1, the orientation on $\mathcal{M}_l(\mathbb{P}^{2m-1},[2d])^{\tau_{2m-1},\eta}$ there still has to be flipped when gluing this space to $\mathcal{M}_l(\mathbb{P}^{2m-1},[2d])^{\tau_{2m-1},\tau}$. Thus, the numbers $N_{2d}^{\phi}(t_1,\ldots,t_l)$ still vanish.

Remark A.6. Let $m\in\mathbb{Z}^+$ be odd so that

$$w_2(\mathbb{RP}^{2m-1})=w_2\big(\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\big)|_{\mathbb{RP}^{2m-1}}.$$

For each $d\in\mathbb{Z}$, denote by

$$[d/2]\in H_2(\mathbb{P}^{2m-1},\mathbb{RP}^{2m-1})\cong\begin{cases}\mathbb{Z}\oplus\mathbb{Z}, & \text{if } n=1;\\ \mathbb{Z}, & \text{if } n\geq 2;\end{cases}$$

a class such that $[d/2]-\tau_{2m-1*}[d/2]$ is the image of $[d]\in H_2(\mathbb{P}^{2m-1})$ under the homomorphism induced by the inclusion $(\mathbb{P}^{2m-1},\emptyset)\to(\mathbb{P}^{2m-1},\mathbb{RP}^{2m-1})$. Under the identification

$$\big(2\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\big)^{\mathbb{R}}\to\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\big|_{\mathbb{RP}^{2m-1}},\qquad(\ell,\alpha_1,\alpha_2)\to(\ell,\alpha_1),\tag{A.21}$$

the complex line bundle $\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\to\mathbb{P}^{2m-1}$, viewed as a real oriented rank 2 vector bundle, and the canonical spin structure on $\mathcal{O}_{\mathbb{P}^{2m-1}}(1)\big|_{\mathbb{RP}^{2m-1}}\oplus T\mathbb{RP}^{2m-1}$, defined as in Section A.2, induce orientations on

$$\mathcal{M}^{\mathrm{disk}}_{0,0}\big(\mathbb{P}^{2m-1},\mathbb{RP}^{2m-1},[d/2]\big)\qquad\text{and}\qquad\mathcal{M}^{\mathrm{disk}}_{0,l}\big(\mathbb{P}^{2m-1},\mathbb{RP}^{2m-1},[d/2]\big)_{\mathrm{dec}};$$

see [8, Section 3.1] and [9, Section 3.3]. By [8, Theorem 1.3] and [8, Section 3.2], the sign of the deck transformation $\tau_{\mathcal{M}}$ for the double cover

$$\mathcal{M}^{\mathrm{disk}}_{0,l}\big(\mathbb{P}^{2m-1},\mathbb{RP}^{2m-1},[d/2]\big)_{\mathrm{dec}}\to\mathcal{M}_l(\mathbb{P}^{2m-1},[d])^{\tau_{2m-1},\tau}$$

with the above orientation on the domain is the parity of

$$\frac{1}{2}\mu\big([d/2]\big)+d=(m+1)d,$$

i.e. $\tau_{\mathcal{M}}$ is orientation-preserving. Remark A.5 implies that this double cover is orientation-preserving with respect to the orientation constructed in Section A.1 on the target. In fact, the argument of Remark A.5 applies without change in the case i is odd, since the pull-back of the trivialization (A.14) by f_0 to

$$S^1\equiv\big\{[x,y]\colon\,|x|\!=\!|y|\big\}\subset D^2\equiv\big\{[x,y]\colon\,|x|\!\le\!|y|\big\}$$

extends to a trivialization of $f_0^*\mathcal{O}_{\mathbb{P}^{2m-1}}(1)$, as an oriented rank 2 vector bundle, over D^2, as required by the construction of [8, Section 3.1]. Since $\tau_{\mathcal{M}}$ is orientation-preserving, this implies that the orientations agree for i even as well. However, this can also be seen directly as follows. We first change Ψ in (A.15) in the i-th factor to

$$(\ell,\alpha_{\bar{i}},\alpha_i)\to\big(\ell,\alpha_{\bar{i}}(\mathbf{z}_i(\ell))\big)=\big(\ell,z_{i\bar{i}}\overline{\alpha_i(\mathbf{z}_i(\ell))}\big);$$

the pull-back of the corresponding trivialization of the rank 2 real vector bundle (A.21) by f_0 to S^1 then extends over D^2. We must thus use this new Ψ to orient $H^0(\Sigma_0,f_0^*T\mathbb{P}^{2m-1})_{\mathbb{R}}$. The components of the change of the trivialization after (A.18) taking the two standard real basis vectors for $\mathbb{C}$ to 1 and $\mathfrak{i}/2a$ now take them to $z_{i\bar{i}}$ and $-z_{i\bar{i}}\mathfrak{i}/2a$; in order for the composite orientation to preserve the orientation, we must thus take $a\!<\!0$. The factor of $z_{i\bar{i}}$ pulls back to z^{d_0} by f_0, giving the same trivialization of $f_0^*T\mathbb{RP}^{2m-1}$ over S^1 if and only if d_0 is even. Thus, the orientation on $H^0(\Sigma_0,f_0^*T\mathbb{P}^{2m-1})_{\mathbb{R}}$ as described above differs from the correct one by $(-1)^{d_0}$. Finally, the characteristic vector field (A.20) is now oriented in the opposite way to the factor of $\mathbb{R}$ in $H^0(\Sigma_0,f_0^*T\mathbb{P}^{2m-1})_{\mathbb{R}}$, which contributes another factor of (-1) to the sign, and an overall sign of $(-1)^{(d_0+1)m\bar{i}}$ to (A.10).

REFERENCES

[1] M. Atiyah and R. Bott, *The moment map and equivariant cohomology*, Topology 23 (1984), 1–28.

[2] M. Aganagic, A. Klemm, and C. Vafa, *Disk instantons, mirror symmetry and the duality web*, Z. Naturforsch. A 57 (2002), no. 1–2, 1–28.

[3] Mohammad. Farajzadeh Tehrani, *Open Gromov-Witten theory on symplectic manifolds via symplectic cutting*, Advances in Mathematics 232 (2013) 238–270.

[4] Mohammad Farajzadeh Tehrani, *On moduli spaces of real curves in symplectic manifolds*, PhD Thesis, 2012, Princeton university.

[5] K. Fukaya, *Counting pseudo-holomorphic discs in Calabi-Yau 3 fold*, Tohoku Math. J. (2) 63 (2011), no. 4, 697-727.

[6] K. Fukaya, and K. Ono, *Arnold conjecture and Gromov-Witten invariant*, Topology 38 (1999), no. 5, 933-1048.

[7] K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono, *Lagrangian intersection Floer theory: anomaly and obstruction, Volumes I,II*, AMS/IP Studies in Advanced Mathematics, 2009.

[8] K. Fukaya, Y-G, Oh. H, Ohta and K. Ono, *Anti-symplectic involution and Floer cohomology*, arXiv:0912.2646.

[9] P. Georgieva, *Orientability of moduli spaces and open Gromov-Witten invariants*, PhD Thesis, 2011.

[10] K. Hori, S. Katz, A. Klemm, R. Pandharipande, R. Thomas, C. Vafa, R. Vakil, E. Zaslow, *Mirror Symmetry*, Clay Mathematics Monographs.

[11] J. M. F. Labastida, M. Marino, and C. Vafa, *Knots, links and branes at large N*, J. High Energy Phys. 2000, no. 11, Paper 7, 42 pp.

[12] J. Li and G. Tian, *Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties*, J. Amer. Math. Soc. 11 (1998), no. 1, 119–174.

[13] D. McDuff and D. Salamon, *J-holomorphic curves and symplectic topology*, American Mathematical Society Colloquium Publications, 52, American Mathematical Society, Providence, RI, 2004.

[14] H. Ooguri and C. Vafa, *Knot invariants and topological strings*, Nuclear Phys. B 577 (2000), no. 3, 419–438.

[15] R. Pandharipande, J. Solomon, and J. Walcher, *Disk enumeration on the quintic 3-fold* J. Amer. Math. Soc. 21 (2008), no. 4, 1169-1209.

[16] A. Popa and A. Zinger, *Mirror Symmetry for Closed, Open, and Unoriented Gromov-Witten Invariants*, arXiv:1010.1946.

[17] Y. Ruan and G. Tian, *A mathematical theory of quantum cohomology*, J.Differential Geom. 42 (1995), no. 2, 259–367.

[18] J. Solomon, *Intersection theory on the moduli space of holomorphic curves with Lagrangian boundary conditions*, arXiv:math/0606429.

[19] – , *Open Gromov-Witten invariants in dimension four*, arXiv:1110.2705.

[20] – , *Open Gromov-Witten invariants in dimension six* arXiv:1201.3518.

[21] E. Witten, *Chern-Simons gauge theory as a string theory*, The Floer memorial volume, 637-678, Progr. Math., 133, Birkhauser, Basel, 1995.

Department of Mathematics, Cornell University
E-mail address: `tehrani@math.cornell.edu`